

2020

Peanut Variety and Quality Evaluation Results

Quality Data

**Tidewater Agricultural Research and
Extension Center**

Virginia Agricultural Experiment Station



**Virginia
Cooperative
Extension**

Virginia Tech
Virginia State University

PEANUT VARIETY AND QUALITY EVALUATION RESULTS 2020

II. Quality Data

Maria Balota, Ph.D.
Associate Professor Crop Physiology
Virginia Tech – Tidewater AREC

Jeffrey Dunne, Ph.D.
Assistant Professor, Peanut Breeder
North Carolina State University

Alexandre Brice Cazenave, Ph.D.
Research Associate
Virginia Tech – Tidewater AREC

Dan Anco, Ph.D.
Peanut Extension Specialist
Clemson University

Wayne Nixon
Agronomist
Severn Peanut Co.

TECHNICAL SUPPORT:

F. Bryant, Ag Specialist
Z. Dunlow, Ag Specialist
F. Cherry, Ag Technician

Virginia Polytechnic Institute and State University
Virginia Agricultural Experiment Station
Tidewater Agricultural Research and Extension Center
Suffolk, Virginia 23437

March 2021

Virginia Tech does not discriminate against employees, students, or applicants on the basis of race, color, sex, disability, age, veteran status, national origin, religion, sexual orientation, or political affiliation. Anyone having questions concerning discrimination or accessibility should contact the Equal Opportunity/Affirmative Action Office.

ACKNOWLEDGEMENTS

FINANCIAL SUPPORT

The authors gratefully acknowledge financial support from the following institutions and organizations:

Virginia Tech

Virginia Agricultural Experiment Station

NC State University

National Peanut Shellers Association

South Carolina Peanut Growers

North Carolina Peanut Growers

Virginia Crop Improvement Association



TECHNICAL SUPPORT

The following agricultural specialists, technicians, and lab assistants are gratefully acknowledged for their professionalism, and dedication to achieve tasks on time and in a collegial manner: Frank Bryant, Fitz Cherry, and Zoe Dunlow.



Zoe Dunlow



Fitz Cherry



Frank Bryant

All of the following cooperators are also acknowledged for their various support provided to the PVQE program in 2020.

LIST OF COOPERATORS

Virginia Tech, Virginia Agricultural Experiment Station, and VCIA

- Mr. K. Jones, Farm Manager, Tidewater AREC
- Mr. B. Slye, Assistant Farm Manager, Tidewater AREC
- Mr. P. Browning, VCIA
- Mr. T. Hardiman, VCIA

Other universities

- Dr. J. Dunne, NCSU
- Dr. D. Anco, Clemson
- Dr. B. Tillman, University of Florida
- Mr. C. Bogle, Upper Coastal Plain Research Station, NCSU

Growers

- Mr. T. Slade, Martin Co., NC
- Mr. D. McDuffie, Bladen Co., NC

County Agents

- Ms. L. Preisser, Isle of Wight Co., VA
- Mr. S. Reiter, Prince George Co., VA
- Mr. M. Parrish, Dinwiddie Co., VA
- Ms. E. Pittman, Suffolk Co. VA
- Ms. S. Rutherford, Greensville Co. VA
- Mr. J. Holland, Southampton Co., VA
- Mr. L. Grimes, Martin Co., NC

Commodity Groups

- Mr. D. Cotton, Virginia Peanut Board
- Mr. B. Sutter, North Carolina Peanut Board
- Ms. M. Copelan, South Carolina Peanut Board

Companies

- Mr. M. Simmons, Birdsong Peanut
- Mr. K. Bennett, Birdsong Peanut
- Mr. J. Laine, Wakefield Peanut Company
- Mr. B. Gwaltney, Indika Farms, Inc.

- | | |
|------------------------------|--------------------------|
| Amadas Industries | Dow Agro Sciences LLC |
| AMVAC | DuPont |
| BASF Corporation | Helena |
| Bayer Crop Science | Monsanto |
| Coastal Chemical Corporation | Syngenta Crop Protection |
| | Valent USA Corporation |

TABLE OF CONTENTS

Acknowledgements.....	i
Technical Support.....	iii
List of Cooperators	iv
Table of Contents.....	v
List of Tables	vi
Introduction.....	1
Plant Material and Test Location.....	2
2020 Blanching Results	4
2020 Fatty Acid Results.....	22

List of Tables

1.	Breeding lines and varieties evaluated in 2020	2
2.	Laboratory sample blanching of Extra Large Kernels (ELK) from Tidewater AREC (Suffolk), VA, Dig 1, 2020	4
3.	Laboratory sample blanching of Extra Large Kernels (ELK) from Tidewater AREC (Suffolk), VA, Dig 2, 2020	5
4.	Laboratory sample blanching of Extra Large Kernels (ELK) from Tidewater AREC (Suffolk), VA, Average of all Digs, 2020	6
5.	Laboratory sample blanching of Extra Large Kernels (ELK) from Martin County, NC, Dig 1, 2020	7
6.	Laboratory sample blanching of Extra Large Kernels (ELK) from Martin County, NC, Dig 2, 2020	8
7.	Laboratory sample blanching of Extra Large Kernels (ELK) from Martin County, NC, Average of all Digs, 2020	9
8.	Laboratory sample blanching of Extra Large Kernels (ELK). Averages from Tidewater AREC (Suffolk), VA and Martin County, NC, 2020	10
9.	Laboratory sample blanching of Extra Large Kernels (ELK). Averages from Tidewater AREC (Suffolk), VA and Martin County, NC., Two-year averages (2019-2020)	11
10.	Laboratory sample blanching of Extra Large Kernels (ELK). Averages from Tidewater AREC (Suffolk), VA and Martin County, NC., Three-year averages (2018-2020)	12
11.	Laboratory sample blanching of Medium Kernels from Tidewater AREC (Suffolk), VA, Dig 1, 2020	13
12.	Laboratory sample blanching of Medium Kernels from Tidewater AREC (Suffolk), VA, Dig 2, 2020	14
13.	Laboratory sample blanching of Medium Kernels from Tidewater AREC (Suffolk), VA, Average of all Digs, 2020	15
14.	Laboratory sample blanching of Medium Kernels from Martin County, NC, Dig 1, 2020	16
15.	Laboratory sample blanching of Medium Kernels from Martin County, NC, Dig 2, 2020	17
16.	Laboratory sample blanching of Medium Kernels from Martin County, NC, Average of all Digs, 2020	18
17.	Laboratory sample blanching of Medium Kernels. Averages from Tidewater AREC (Suffolk), VA and Martin County, NC, 2020	19
18.	Laboratory sample blanching of Medium Kernels. Averages from Tidewater AREC (Suffolk), VA and Martin County, NC, Two-year averages (2019-2020)	20
19.	Laboratory sample blanching of Medium Kernels. Averages from Tidewater AREC (Suffolk), VA and Martin County, NC., Three-years averages (2018-2020)	21

List of tables continued

20.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk), VA, Dig 1, 2020.....	22
21.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk), VA, Dig 2, 2020.....	24
22.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk), VA, average of all Digs, 2020	26
23.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC, Dig 1, 2020.....	28
24.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC, Dig 2, 2020.....	30
25.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC, Average of all Dig, 2020.....	32
26.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Rocky Mount, NC, 2020	34
27.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Bladen, NC, 2020.....	36
28.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Blackville, SC, 2020	38
29.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Averaged across all locations, 2020	40
30.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Two-year averages across all locations, (2019- 2020).....	42
31.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Three-year averages across all locations, (2018- 2020).....	44
32.	Fatty Acid Composition, Iodine Values, Oleic/Linoleic O/L Ratio, %Total Saturated, Total Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Rain Shelter Trial, Suffolk, VA, 2020	46

33. Fatty Acid Composition, Iodine Values, Oleic/Linoleic O/L Ratio, %Total Saturated,
Total Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated.
Rain Shelter Trial, Suffolk, VA, (2019 – 2020) Two-year averages48

INTRODUCTION

Along with agronomic and grade information, data on kernel and pod quality are essential for release of new peanut cultivars to ensure acceptability by the entire peanut trade. The present report contains the quality data collected on 6 Virginia-type cultivars that currently are on the market and 24 advanced breeding lines tested in the Peanut Variety and Quality Evaluation (PVQE) small plots in 2020. The small PVQE plots with 30 varieties were tested at five locations in Virginia, North Carolina, and South Carolina: Suffolk, VA, Martin Co., NC, Rocky Mount, NC, Bladen Co., NC, and Blackville, SC. At Suffolk, VA and at Martin Co., NC, two digs were achieved. For the other locations, only one dig was tested. Each genotype was replicated 2 times at each location and digging date. Varieties' names and pedigree are presented in Table 1. A detailed description of the plant material, test locations, weather conditions, and cultural practices is included in the PVQE 2020 Results. I. Agronomic and Grade Data, at [SPES-181.pdf \(vt.edu\)](https://www.vt.edu/SPES-181.pdf).

2020 SMALL PLOT TESTS

Blanching evaluations were determined by a laboratory sample blancher of two, 250 g peanut samples from two dig dates at Martin Co., NC, and the Tidewater AREC. Tables 2 through 19 contain blanching data for the extra-large kernels (ELK) and medium-size kernels. Statistical analyses were determined for percentage of splits, whole blanched, not blanched, and partially blanched.

Small Plot Tests

PLANT MATERIAL AND TEST LOCATIONS

Table 1. Names and pedigree of the genotypes (advanced breeding lines and commercial varieties) evaluated in 2020.

Genotype number	Variety/line	Parentage
1	Bailey	NC 12C*2 / N96076L
2	Bailey II	Bailey / XO7016 (BC2F1 – 04:F01)
3	Emery	N03079FT*2 / Brantley
4	Sullivan	Bailey / X03034 (F01)
5	Walton	2000x10-1-B2-3-2-2/97x48-HO3-7-B2-2-b3-B
6	Wynne	N03079FT / X03034 (F01)
7	N14001	N02006 // X05012, N02006 / N02064ol
8	N14002olJ	N03079FT // X05024, N03079FT / N02064ol
9	N14004olJ	Bailey // X05027, Bailey / N02060ol
10	N14007	Phillips / N99121CSm, X00044 /3/ X05036, Phillips / N99121CSm, X00044 // N02064ol
11	N14009	Phillips / N99121CSm, X00044 /3/ X05036, Phillips / N99121CSm, X00044 // N02064ol
12	N15066	N02054ol // N02005 / N02054ol, X03138 /3/ N03084FT
13	N14023ol ¹	N01015T / N00098ol, X02083 // Sugg
14	N14027olJ	Bailey /4/ X07019, Bailey // X05028, Bailey / N02064ol, X05250 /3/ Bailey
15	N15017ol	Bailey /4/ X07018, Bailey // X05028, Bailey / N02064ol, X05250 /3/ Bailey
16	N15039ol	N03079FT*2 / N02054ol, X03153 // N05042F
17	N15041ol	N03079FT*2 / N02059ol, X03155 // N05044FCSm
18	N15044olF	N03079FT*2 / N02059ol, X03155 // N05044FCSm
19	N15053	N08082olJCT // X09019, N08082olJCT / Florida Fancy
20	N15060	Bailey*2 / Brantley, N08086olJCT // SPT 07-01, NC-V 11 / GP-NC WS 11
21	N16005	Bailey*2 / Brantley, X03157 // GP-NC WS 16
22	N16012	N08082olJCT /3/ X09008, N08082olJCT // SPT 07-01, NC-V 11 / GP-NC WS 11
23	N16021	N08082olJCT // X09019, N08082olJCT / Florida Fancy
24	N17036	Emery /3/ N11035olSrT, N03079FT*2 / Brantley, X03151 // Sugg
25	N17037	Emery /3/ N11035olSrT, N03079FT*2 / Brantley, X03151 // Sugg
26	N17040	N03079FT*2 / Brantley, N10047ol // N12010ol, Bailey*4 / N02060ol
27	N17041	N03079FT*2 / Brantley, N10047ol // N12010ol, Bailey*4 / N02060ol
28	N17044	Bailey*2 / Brantley, N10053ol // Bailey II, Bailey*4 / N02060ol
29	N17045	Bailey*2 / Brantley, N10053ol /3/ CRSP 1050-110, Florida MDR 98 / Bayo Grande, 0020-20 // FNC94022-1-2-1-1-b3-B , N91026E / PI 576638
30	N17047	Bailey*2 / Brantley, N10053ol /3/ CRSP 1050-110, Florida MDR 98 / Bayo Grande, 0020-20 // FNC94022-1-2-1-1-b3-B , N91026E / PI 576638

¹ N14023ol was released as a cultivar in 2020, as 'NC 20'.

Small Plot Tests

Fatty acid content and composition of the sound mature kernels (SMK) was determined by gas chromatography and expressed as % from total seed oil content. Iodine value, oleic/linoleic (O/L) ratio, % total saturated, polyunsaturated/saturated (P/S) ratio, and % total long chain-saturated acids were also calculated using the following formulas:

$$\text{Iodine Value} = (\% \text{ oleic}) (0.8601) + (\% \text{ linoleic}) (1.7321) + (\% \text{ eicosenoic}) (0.7854)$$

$$\text{Oleic/Linoleic (O/L) ratio} = \% \text{ oleic} / \% \text{ linoleic}$$

$$\% \text{ Total Saturated} = \% \text{ palmitic} + \% \text{ stearic} + \% \text{ arachidic} + \% \text{ behenic} + \% \text{ lignoceric}$$

$$\text{Polyunsaturated/Saturated (P/S) ratio} = \% \text{ polyunsaturated (linoleic)} / \% \text{ total saturated}$$

$$\% \text{ Total Long Chain Saturated} = \% \text{ arachidic} + \% \text{ behenic} + \% \text{ lignoceric}$$

The definition of a high oleic peanut is a peanut line and seed that has an oleic acid content of from about 74% to about 84% and a linoleic acid content of from about 2% to about 8%, each based upon the total fatty acid content of the seed, and a ratio of the amount of oleic acid to linoleic acid in the seed of from about 9:1 to about 42:1.

Fatty acid composition is reported from all 2020 PVQE locations and digging dates in Tables 20 through 28. Table 29 shows the content of the fatty acids averaged across all locations in 2020. Two- and three-year averages are included in Tables 30 and 31. Finally, fatty acid composition of 14 cultivars and breeding lines exposed to soil water deficit stress is reported in Tables 32 and 33.

Statistics:

Analysis of Variance was run for the cultivars and breeding lines on individual digging dates and locations, and averaged digging dates, locations, and years. When significant differences between cultivars and lines were detected, means were compared by the Fisher's LSD test and the minimum significant difference was included in the tables. These values were used to compare the varieties.

Blanching Results

Table 2. Laboratory sample blanching of Extra-Large Kernels (ELK) from Tidewater AREC (Suffolk, VA), Dig 1, 2020 (28 September).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.6 a-c	4.7 a	0.5 b	1.2 e	94.7 ab	0.0 b	2.5 fg
Bailey II	5.6 bc	4.7 a	0.9 a	1.7 b-e	94.7 a	0.0 b	2.0 g
Emery	5.7 ab	4.8 a	0.9 a	1.2 e	94.5 a-c	0.0 b	2.7 e-g
Sullivan	5.6 a-c	4.8 a	0.8 a	1.3 de	93.9 a-d	0.0 b	3.3 c-g
Walton	5.6 bc	4.7 a	0.9 a	1.5 c-e	92.3 de	0.0 b	4.6 a-c
Wynne	5.6 bc	4.8 a	0.8 a	1.3 de	92.8 b-e	0.0 b	4.3 b-e
N14001	5.6 bc	4.7 a	0.9 a	2.0 a-e	91.9 e	0.0 b	4.6 a-c
N14002oIJ	5.6 a-c	4.8 a	0.9 a	2.6 a-e	91.9 e	0.0 b	3.9 b-f
N14004oIJ	5.6 a-c	4.7 a	0.9 a	2.0 a-e	91.9 e	0.0 b	4.5 b-d
N14007	5.6 bc	4.8 a	0.8 a	2.2 a-e	89.9 f	0.0 b	6.2 a
N14009	5.7 ab	4.8 a	0.9 a	2.6 a-d	92.6 c-e	0.0 b	3.1 c-g
N15066	5.5 c	4.7 a	0.8 a	3.4 a	89.7 f	0.2 a	5.1 ab
N14023oI	5.7 ab	4.8 a	0.9 a	1.9 b-e	92.9 a-e	0.0 b	3.6 b-g
N14027oIJ	5.6 a-c	4.8 a	0.8 a	2.2 a-e	92.6 de	0.2 a	3.5 b-g
N15017oI	5.6 a-c	4.7 a	0.9 a	2.0 a-e	92.7 c-e	0.0 b	3.6 b-g
N15039oI	5.7 a	4.8 a	0.9 a	2.9 a-c	92.5 de	0.0 b	3.0 c-g
N15041oI	5.6 a-c	4.8 a	0.9 a	1.2 de	93.7 a-e	0.0 b	3.6 b-g
N15044oIF	5.6 a-c	4.8 a	0.8 a	2.2 a-e	93.0 a-e	0.0 b	3.2 c-g
N15053	5.7 ab	4.8 a	0.9 a	1.7 b-e	93.4 a-e	0.0 b	3.3 c-g
N15060	5.6 a-c	4.8 a	0.9 a	2.6 a-d	92.5 de	0.2 ab	3.1 c-g
N16005	5.7 ab	4.8 a	0.9 a	1.9 b-e	93.9 a-d	0.0 b	2.7 e-g
N16012	5.5 c	4.7 a	0.8 a	2.3 a-e	93.2 a-e	0.0 b	2.8 d-g
N16021	5.6 a-c	4.7 a	0.9 a	2.5 a-e	91.9 e	0.0 b	4.0 b-f
N17036	5.6 bc	4.8 a	0.8 a	2.5 a-e	92.8 a-e	0.0 b	3.1 c-g
N17037	5.7 ab	4.8 a	0.9 a	2.3 a-e	92.6 c-e	0.0 b	3.4 b-g
N17040	5.6 bc	4.8 a	0.8 a	2.3 a-e	93.0 a-e	0.0 b	3.1 c-g
N17041	5.6 bc	4.8 a	0.8 a	3.0 ab	92.8 b-e	0.0 b	2.6 e-g
N17044	5.5 c	4.7 a	0.8 a	2.5 a-e	92.5 de	0.0 b	3.4 b-g
N17045	5.6 a-c	4.7 a	0.9 a	2.1 a-e	92.7 c-e	0.0 b	3.6 b-g
N17047	5.6 bc	4.8 a	0.8 a	1.9 b-e	93.6 a-e	0.0 b	2.9 c-g
Mean	5.6	4.7	0.8	2.1	92.7	0.0	3.5
LSD¹	0.1	0.1	0.3	1.4	1.9	0.2	1.7

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 3. Laboratory sample blanching of Extra-Large Kernels (ELK) from Tidewater AREC (Suffolk, VA), Dig 2, 2020 (20 October).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.5 b	4.7 a	0.8 ab	2.9 a-c	92.0 a-f	0.0 a	3.4 c-g
Bailey II	5.6 ab	4.8 a	0.8 b	1.8 c	92.0 a-f	0.0 a	4.6 b-d
Emery	5.5 b	4.7 a	0.8 ab	2.3 a-c	93.1 a-c	0.0 a	3.0 e-g
Sullivan	5.5 b	4.7 a	0.8 ab	2.4 a-c	92.5 a-f	0.0 a	3.5 c-g
Walton	5.6 a	4.8 a	0.9 ab	2.3 a-c	91.9 a-f	0.2 a	4.1 b-g
Wynne	5.5 b	4.8 a	0.8 b	2.7 a-c	90.9 d-g	0.2 a	4.6 b-d
N14001	5.6 ab	4.8 a	0.8 ab	3.0 a-c	91.1 b-f	0.0 a	4.3 b-f
N14002oIJ	5.6 a	4.8 a	0.9 ab	2.8 a-c	92.2 a-f	0.0 a	3.3 d-g
N14004oIJ	5.6 ab	4.8 a	0.8 ab	2.9 a-c	91.8 a-f	0.2 a	3.6 b-g
N14007	5.5 b	4.7 a	0.8 ab	2.5 a-c	89.0 g	0.2 a	6.7 a
N14009	5.6 a	4.8 a	0.9 ab	2.4 a-c	91.4 a-f	0.2 a	4.5 b-e
N15066	5.6 ab	4.8 a	0.8 ab	2.3 a-c	92.7 a-d	0.2 a	3.2 d-g
N14023oI	5.5 b	4.8 a	0.8 b	2.3 a-c	91.3 a-f	0.2 a	4.6 b-d
N14027oIJ	5.6 a	4.8 a	0.9 ab	2.1 a-c	91.4 a-f	0.0 a	4.9 bc
N15017oI	5.6 ab	4.8 a	0.8 b	2.9 a-c	91.9 a-f	0.0 a	3.6 b-g
N15039oI	5.6 ab	4.8 a	0.8 ab	3.1 a-c	90.7 e-g	0.0 a	4.7 b-d
N15041oI	5.5 b	4.8 a	0.8 b	2.7 a-c	91.6 a-f	0.0 a	4.1 b-g
N15044oIF	5.6 a	4.8 a	0.8 ab	2.6 a-c	90.6 fg	0.2 a	5.0 b
N15053	5.6 a	4.8 a	0.9 ab	2.2 a-c	93.1 ab	0.2 a	2.9 fg
N15060	5.5 b	4.7 a	0.8 ab	2.4 a-c	90.9 d-g	0.2 a	4.9 bc
N16005	5.5 b	4.7 a	0.8 ab	2.7 a-c	91.5 a-f	0.0 a	4.2 b-f
N16012	5.6 ab	4.8 a	0.8 ab	3.0 a-c	92.0 a-f	0.0 a	3.4 d-g
N16021	5.6 ab	4.8 a	0.8 ab	3.4 a	91.1 c-f	0.2 a	3.7 b-g
N17036	5.6 ab	4.8 a	0.8 ab	1.9 c	93.3 a	0.2 a	2.6 g
N17037	5.6 a	4.8 a	0.8 ab	2.0 bc	92.6 a-e	0.2 a	3.6 b-g
N17040	5.6 ab	4.8 a	0.8 ab	1.8 c	92.5 a-f	0.0 a	4.0 b-g
N17041	5.6 ab	4.8 a	0.8 ab	2.6 a-c	91.7 a-f	0.0 a	4.0 b-g
N17044	5.6 ab	4.7 a	0.9 ab	3.3 ab	92.1 a-f	0.0 a	3.0 e-g
N17045	5.6 a	4.7 a	0.9 a	2.3 a-c	92.8 a-d	0.0 a	3.4 d-g
N17047	5.6 a	4.8 a	0.8 ab	2.9 a-c	92.3 a-f	0.0 a	3.2 d-g
Mean	5.6	4.7	0.8	2.5	91.8	0.1	3.9
LSD¹	0.1	0.1	0.1	1.3	2.0	0.3	1.5

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 4. Laboratory sample blanching of Extra-Large Kernels (ELK). Averages of both digging dates from Tidewater AREC (Suffolk, VA), 2020.

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.6 ab	4.7 c	0.6 b	2.0 a-d	93.3 a-c	0.0 b	3.0 de
Bailey II	5.6 ab	4.8 a-c	0.8 a	1.7 d	93.4 ab	0.0 b	3.3 b-e
Emery	5.6 ab	4.7 bc	0.9 a	1.7 d	93.8 a	0.0 b	2.8 e
Sullivan	5.6 ab	4.8 a-c	0.8 a	1.8 cd	93.2 a-d	0.0 b	3.4 b-e
Walton	5.6 ab	4.7 bc	0.9 a	1.9 b-d	92.1 b-f	0.1 ab	4.3 bc
Wynne	5.5 b	4.8 a-c	0.8 a	2.0 a-d	91.8 b-f	0.1 ab	4.4 b
N14001	5.6 ab	4.7 bc	0.8 a	2.5 a-d	91.5 ef	0.0 b	4.4 b
N14002oIJ	5.6 ab	4.8 a-c	0.9 a	2.7 a-d	92.1 b-f	0.0 b	3.6 b-e
N14004oIJ	5.6 ab	4.7 bc	0.9 a	2.4 a-d	91.8 b-f	0.1 ab	4.0 b-e
N14007	5.5 b	4.7 bc	0.8 a	2.4 a-d	89.4 g	0.1 ab	6.5 a
N14009	5.6 a	4.8 a-c	0.9 a	2.5 a-d	92.0 b-f	0.1 ab	3.8 b-e
N15066	5.5 b	4.7 bc	0.8 a	2.8 ab	91.2 f	0.2 a	4.1 b-d
N14023oI	5.6 ab	4.8 a-c	0.8 a	2.1 a-d	92.1 b-f	0.1 ab	4.1 b-d
N14027oIJ	5.6 ab	4.8 ab	0.8 a	2.1 a-d	92.0 b-f	0.1 ab	4.2 b-d
N15017oI	5.6 ab	4.8 a-c	0.8 a	2.4 a-d	92.3 a-f	0.0 b	3.6 b-e
N15039oI	5.6 a	4.8 ab	0.9 a	3.0 a	91.6 d-f	0.0 b	3.8 b-e
N15041oI	5.6 ab	4.8 a-c	0.8 a	2.0 a-d	92.6 a-f	0.0 b	3.8 b-e
N15044oIF	5.6 ab	4.8 a	0.8 a	2.4 a-d	91.8 b-f	0.1 ab	4.1 b-d
N15053	5.6 a	4.8 a-c	0.9 a	1.9 b-d	93.3 a-c	0.1 ab	3.1 c-e
N15060	5.6 ab	4.7 bc	0.8 a	2.5 a-d	91.7 c-f	0.2 ab	4.0 b-e
N16005	5.6 ab	4.8 a-c	0.8 a	2.3 a-d	92.7 a-f	0.0 b	3.4 b-e
N16012	5.5 b	4.7 bc	0.8 a	2.7 a-d	92.6 a-f	0.0 b	3.1 c-e
N16021	5.6 ab	4.7 bc	0.9 a	3.0 a	91.5 ef	0.1 ab	3.8 b-e
N17036	5.6 ab	4.8 a-c	0.8 a	2.2 a-d	93.0 a-c	0.1 ab	2.8 e
N17037	5.6 a	4.8 ab	0.9 a	2.2 a-d	92.6 a-f	0.1 ab	3.5 b-e
N17040	5.6 ab	4.8 a-c	0.8 a	2.1 a-d	92.7 a-f	0.0 b	3.6 b-e
N17041	5.6 ab	4.8 a-c	0.8 a	2.8 a-c	92.2 a-f	0.0 b	3.3 b-e
N17044	5.5 b	4.7 c	0.8 a	2.9 ab	92.3 a-f	0.0 b	3.2 b-e
N17045	5.6 ab	4.7 c	0.9 a	2.2 a-d	92.7 a-f	0.0 b	3.5 b-e
N17047	5.6 ab	4.8 a	0.8 a	2.4 a-d	92.9 a-e	0.0 b	3.0 de
Mean	5.6	4.7	0.8	2.3	92.3	0.0	3.7
LSD¹	0.1	0.1	0.1	1.0	1.6	0.2	1.3

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 5. Laboratory sample blanching of Extra-Large Kernels (ELK) from Martin County, NC, Dig 1, 2020 (5 October).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 a	4.8 a	0.9 a	1.3 e	94.4 a-c	0.0 a	2.7 bc
Bailey II	5.6 ab	4.8 a	0.8 cd	1.3 e	94.9 a	0.0 a	2.1 c
Emery	5.6 ab	4.8 a	0.8 bc	1.5 c-e	94.8 ab	0.0 a	2.1 c
Sullivan	5.6 ab	4.8 a	0.8 bc	2.0 b-e	92.4 b-d	0.0 a	4.0 bc
Walton	5.6 ab	4.8 a	0.8 bc	2.2 b-e	92.5 b-d	0.0 a	3.7 bc
Wynne	5.6 ab	4.8 a	0.9 ab	2.2 b-e	91.8 d	0.2 a	4.2 bc
N14001	5.6 ab	4.8 a	0.8 bc	2.1 b-e	92.4 cd	0.0 a	3.9 bc
N14002oIJ	5.6 ab	4.8 a	0.8 bc	2.3 a-e	92.8 a-d	0.0 a	3.3 bc
N14004oIJ	5.6 ab	4.8 a	0.9 ab	2.2 b-e	92.0 d	0.2 a	4.0 bc
N14007	5.5 b	4.8 a	0.7 d	2.7 a-d	88.2 e	0.2 a	7.3 a
N14009	5.6 ab	4.8 a	0.8 bc	1.8 b-e	92.6 a-d	0.0 a	4.0 bc
N15066	5.6 ab	4.8 a	0.9 ab	2.3 a-e	91.7 d	0.0 a	4.4 bc
N14023oI	5.6 ab	4.8 a	0.8 bc	3.0 ab	92.3 cd	0.0 a	3.1 bc
N14027oIJ	5.6 ab	4.8 a	0.8 bc	2.1 b-e	92.4 cd	0.0 a	3.9 bc
N15017oI	5.7 a	4.8 a	0.9 ab	2.4 a-e	92.3 cd	0.0 a	3.8 bc
N15039oI	5.7 a	4.8 a	0.9 ab	2.4 a-e	93.0 a-d	0.0 a	3.0 bc
N15041oI	5.6 ab	4.8 a	0.8 bc	1.5 de	92.7 a-d	0.2 a	4.0 bc
N15044oIF	5.6 ab	4.8 a	0.8 bc	2.3 a-e	90.9 d	0.2 a	5.2 ab
N15053	5.6 ab	4.8 a	0.8 bc	2.0 b-e	91.8 d	0.0 a	4.5 bc
N15060	5.6 ab	4.8 a	0.8 bc	2.2 b-e	93.2 a-d	0.0 a	2.9 bc
N16005	5.6 ab	4.8 a	0.8 bc	2.0 b-e	92.6 a-d	0.0 a	3.8 bc
N16012	5.6 ab	4.8 a	0.9 ab	2.2 b-e	92.7 a-d	0.2 a	3.4 bc
N16021	5.5 b	4.7 a	0.8 bc	3.0 ab	92.1 cd	0.0 a	3.3 bc
N17036	5.6 ab	4.8 a	0.8 bc	3.0 ab	92.4 b-d	0.0 a	2.9 bc
N17037	5.6 ab	4.8 a	0.8 bc	2.2 b-e	93.1 a-d	0.2 a	2.9 bc
N17040	5.5 b	4.7 a	0.8 bc	3.0 ab	92.5 b-d	0.0 a	2.9 bc
N17041	5.6 ab	4.8 a	0.8 bc	2.8 a-c	92.6 a-d	0.0 a	3.0 bc
N17044	5.6 ab	4.8 a	0.8 cd	3.5 a	91.6 d	0.0 a	3.3 bc
N17045	5.6 ab	4.8 a	0.8 bc	2.6 a-d	92.6 a-d	0.0 a	3.1 bc
N17047	5.6 ab	4.8 a	0.8 bc	2.1 b-e	92.5 b-d	0.2 a	3.6 bc
Mean	5.6	4.8	0.8	2.3	92.4	0.0	3.6
LSD¹	0.1	0.1	0.1	1.3	2.4	0.2	2.5

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 6. Laboratory sample blanching of Extra-Large Kernels (ELK) from Martin County, NC, Dig 2, 2020 (15 October).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 ab	4.8 b	0.9 a	2.1 a	92.8 ab	0.0 a	3.5 c-g
Bailey II	5.6 ab	4.8 b	0.9 ab	1.7 a	93.9 a	0.0 a	2.8 fg
Emery	5.6 ab	4.8 b	0.9 ab	1.8 a	93.9 a	0.0 a	2.7 g
Sullivan	5.6 b	4.8 b	0.8 ab	1.9 a	92.4 a-c	0.0 a	4.1 b-g
Walton	5.7 ab	4.8 ab	0.9 ab	2.7 a	91.7 a-d	0.0 a	4.0 b-g
Wynne	5.6 ab	4.8 b	0.9 ab	2.1 a	91.3 b-e	0.2 a	4.8 b-d
N14001	5.6 ab	4.8 ab	0.8 ab	2.0 a	92.2 a-c	0.2 a	4.0 b-g
N14002oIJ	5.6 ab	4.9 a	0.8 b	2.6 a	91.5 a-e	0.0 a	4.2 b-g
N14004oIJ	5.6 ab	4.8 ab	0.8 ab	2.4 a	92.2 a-c	0.0 a	3.8 c-g
N14007	5.6 ab	4.8 ab	0.8 ab	3.2 a	89.3 d-f	0.3 a	5.6 ab
N14009	5.6 ab	4.8 ab	0.8 ab	2.8 a	90.2 c-f	0.2 a	5.2 bc
N15066	5.6 b	4.8 b	0.8 ab	2.2 a	93.2 ab	0.0 a	3.0 e-g
N14023oI	5.6 ab	4.8 b	0.9 ab	3.3 a	89.2 ef	0.4 a	5.6 ab
N14027oIJ	5.7 ab	4.8 ab	0.9 ab	3.0 a	91.2 b-e	0.0 a	4.2 b-g
N15017oI	5.6 b	4.8 b	0.8 ab	2.6 a	91.1 b-e	0.2 a	4.6 b-f
N15039oI	5.6 ab	4.8 ab	0.8 ab	3.1 a	91.1 b-e	0.0 a	4.3 b-g
N15041oI	5.6 ab	4.8 b	0.9 ab	2.6 a	88.5 f	0.3 a	7.1 a
N15044oIF	5.6 ab	4.8 ab	0.8 ab	2.7 a	90.3 c-f	0.2 a	5.2 bc
N15053	5.6 ab	4.8 ab	0.8 ab	3.2 a	91.2 b-e	0.4 a	3.6 c-g
N15060	5.6 ab	4.8 ab	0.8 ab	3.3 a	90.3 c-f	0.4 a	4.4 b-g
N16005	5.6 ab	4.8 ab	0.8 ab	3.1 a	92.0 a-c	0.2 a	3.2 d-g
N16012	5.6 b	4.8 ab	0.8 b	2.3 a	92.0 a-c	0.0 a	4.1 b-g
N16021	5.7 a	4.8 ab	0.9 a	1.8 a	92.0 a-c	0.0 a	4.7 b-e
N17036	5.7 ab	4.9 a	0.8 ab	2.9 a	91.1 b-e	0.3 a	4.1 b-g
N17037	5.6 ab	4.8 ab	0.8 ab	2.4 a	92.8 ab	0.2 a	2.8 fg
N17040	5.6 b	4.8 b	0.8 ab	2.6 a	91.6 a-e	0.0 a	4.2 b-g
N17041	5.6 ab	4.8 ab	0.8 ab	2.6 a	92.3 a-c	0.2 a	3.3 d-g
N17044	5.6 ab	4.8 ab	0.8 ab	2.9 a	91.9 a-c	0.0 a	3.6 c-g
N17045	5.7 ab	4.9 a	0.8 ab	1.9 a	92.5 a-c	0.2 a	3.9 b-g
N17047	5.6 ab	4.8 ab	0.8 ab	2.5 a	92.9 ab	0.2 a	2.9 fg
Mean	5.6	4.8	0.8	2.5	91.6	0.1	4.1
LSD¹	0.1	0.1	0.1	1.7	2.5	0.4	1.8

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 7. Laboratory sample blanching of Extra-Large Kernels (ELK). Averages of both digging dates from Martin County, NC, 2020.

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 a	4.8 ab	0.9 a	1.7 d-f	93.6 ab	0.0 b	3.1 d-g
Bailey II	5.6 a-c	4.8 ab	0.8 b-d	1.5 f	94.4 a	0.0 b	2.5 fg
Emery	5.6 a-c	4.8 ab	0.8 bc	1.6 ef	94.3 a	0.0 b	2.4 g
Sullivan	5.6 bc	4.8 ab	0.8 b-d	1.9 c-f	92.4 b-d	0.0 b	4.0 c-e
Walton	5.6 a-c	4.8 ab	0.8 bc	2.4 a-f	92.1 b-e	0.0 b	3.8 c-g
Wynne	5.6 a-c	4.8 ab	0.9 ab	2.1 b-f	91.5 c-e	0.2 ab	4.5 b-d
N14001	5.6 a-c	4.8 ab	0.8 b-d	2.0 b-f	92.3 b-e	0.1 ab	4.0 c-f
N14002oIJ	5.6 a-c	4.8 a	0.8 cd	2.5 a-e	92.1 b-e	0.0 b	3.8 c-g
N14004oIJ	5.6 a-c	4.8 ab	0.8 bc	2.3 a-f	92.1 b-e	0.1 ab	3.9 c-g
N14007	5.6 bc	4.8 a	0.8 d	2.9 ab	88.8 f	0.2 ab	6.4 a
N14009	5.6 a-c	4.8 ab	0.8 b-d	2.3 a-f	91.4 c-e	0.1 ab	4.6 b-d
N15066	5.6 a-c	4.8 ab	0.8 bc	2.2 a-f	92.4 b-d	0.0 b	3.7 c-g
N14023oI	5.6 a-c	4.8 ab	0.8 bc	3.1 a	90.7 de	0.2 ab	4.3 b-e
N14027oIJ	5.6 a-c	4.8 ab	0.8 bc	2.6 a-e	91.8 c-e	0.0 b	4.1 b-e
N15017oI	5.6 a-c	4.8 ab	0.8 bc	2.5 a-e	91.7 c-e	0.1 ab	4.2 b-e
N15039oI	5.6 ab	4.8 a	0.8 bc	2.7 a-c	92.0 b-e	0.0 b	3.6 d-g
N15041oI	5.6 a-c	4.8 ab	0.8 bc	2.0 b-f	90.6 e	0.2 a	5.5 ab
N15044oIF	5.6 a-c	4.8 a	0.8 b-d	2.5 a-e	90.6 e	0.2 ab	5.2 a-c
N15053	5.6 a-c	4.8 ab	0.8 b-d	2.6 a-d	91.5 c-e	0.2 ab	4.1 b-e
N15060	5.6 a-c	4.8 a	0.8 b-d	2.7 a-c	91.8 c-e	0.2 ab	3.7 c-g
N16005	5.6 a-c	4.8 ab	0.8 b-d	2.5 a-e	92.3 b-e	0.1 ab	3.5 d-g
N16012	5.6 a-c	4.8 ab	0.8 b-d	2.2 a-f	92.3 b-e	0.1 ab	3.7 c-g
N16021	5.6 a-c	4.8 ab	0.9 ab	2.4 a-f	92.0 b-e	0.0 b	4.0 c-f
N17036	5.6 a-c	4.8 a	0.8 b-d	3.0 ab	91.7 c-e	0.2 ab	3.5 d-g
N17037	5.6 a-c	4.8 ab	0.8 b-d	2.3 a-f	92.9 a-c	0.2 ab	2.8 e-g
N17040	5.5 c	4.7 b	0.8 b-d	2.8 a-c	92.0 b-e	0.0 b	3.6 d-g
N17041	5.6 a-c	4.8 ab	0.8 b-d	2.7 a-c	92.4 b-d	0.1 ab	3.1 d-g
N17044	5.6 a-c	4.8 a	0.8 cd	3.2 a	91.7 c-e	0.0 b	3.4 d-g
N17045	5.6 a-c	4.8 a	0.8 b-d	2.2 a-f	92.5 bc	0.1 ab	3.5 d-g
N17047	5.6 a-c	4.8 ab	0.8 b-d	2.3 a-f	92.7 a-c	0.2 ab	3.2 d-g
Mean	5.6	4.8	0.8	2.4	92.0	0.1	3.8
LSD¹	0.1	0.1	0.1	1.0	1.8	0.2	1.5

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 8. Laboratory sample blanching of Extra-Large Kernels (ELK). Averages from Tidewater AREC (Suffolk, VA) and Martin County, NC, 2020.

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.6 ab	4.7 c	0.8 b	1.9 e-g	93.5 a-c	0.0 b	3.0 h-j
Bailey II	5.6 bc	4.8 a-c	0.8 ab	1.6 g	93.9 ab	0.0 b	2.9 ij
Emery	5.6 a-c	4.7 bc	0.8 ab	1.7 fg	94.1 a	0.0 b	2.6 j
Sullivan	5.6 bc	4.8 a-c	0.8 ab	1.9 e-g	92.8 b-d	0.0 b	3.7 c-i
Walton	5.6 a-c	4.8 a-c	0.8 ab	2.2 c-g	92.1 d-g	0.0 ab	4.1 b-g
Wynne	5.6 bc	4.8 a-c	0.8 ab	2.1 c-g	91.7 d-g	0.1 a	4.5 b-d
N14001	5.6 bc	4.8 a-c	0.8 ab	2.2 b-g	91.9 d-g	0.1 ab	4.2 b-e
N14002oIJ	5.6 a-c	4.8 a-c	0.8 ab	2.6 a-d	92.1 d-g	0.0 b	3.7 c-i
N14004oIJ	5.6 a-c	4.8 a-c	0.8 ab	2.4 a-e	92.0 d-g	0.1 ab	3.9 b-h
N14007	5.5 c	4.8 a-c	0.8 ab	2.6 a-d	89.1 h	0.1 a	6.4 a
N14009	5.6 ab	4.8 a-c	0.8 ab	2.4 a-e	91.7 d-g	0.1 ab	4.2 b-e
N15066	5.6 bc	4.7 bc	0.8 ab	2.5 a-e	91.8 d-g	0.1 ab	3.9 b-h
N14023oI	5.6 a-c	4.8 a-c	0.8 ab	2.6 a-d	91.4 fg	0.1 ab	4.2 b-e
N14027oIJ	5.6 ab	4.8 a-c	0.8 ab	2.3 b-f	91.9 d-g	0.1 ab	4.1 b-f
N15017oI	5.6 a-c	4.8 a-c	0.8 ab	2.4 a-e	92.0 d-g	0.0 ab	3.9 b-h
N15039oI	5.6 a	4.8 ab	0.8 ab	2.8 ab	91.8 d-g	0.0 b	3.7 b-i
N15041oI	5.6 a-c	4.8 a-c	0.8 ab	2.0 d-g	91.6 e-g	0.1 ab	4.7 b
N15044oIF	5.6 ab	4.8 a	0.8 ab	2.5 a-e	91.2 g	0.1 ab	4.6 bc
N15053	5.6 ab	4.8 a-c	0.8 ab	2.3 b-g	92.4 c-f	0.1 ab	3.6 d-i
N15060	5.6 a-c	4.8 a-c	0.8 ab	2.6 a-d	91.7 d-g	0.2 a	3.8 b-h
N16005	5.6 a-c	4.8 a-c	0.8 ab	2.4 a-e	92.5 c-f	0.1 ab	3.4 e-j
N16012	5.6 bc	4.8 a-c	0.8 ab	2.4 a-e	92.5 c-f	0.0 ab	3.4 e-j
N16021	5.6 a-c	4.7 bc	0.9 a	2.7 a-d	91.7 d-g	0.1 ab	3.9 b-h
N17036	5.6 a-c	4.8 a-c	0.8 ab	2.6 a-d	92.4 c-f	0.1 ab	3.1 g-j
N17037	5.6 ab	4.8 a-c	0.8 ab	2.2 b-g	92.8 b-e	0.1 ab	3.2 g-j
N17040	5.5 c	4.7 bc	0.8 ab	2.4 a-e	92.4 c-f	0.0 b	3.6 d-j
N17041	5.6 bc	4.8 a-c	0.8 ab	2.7 a-c	92.3 c-g	0.1 ab	3.2 f-j
N17044	5.6 bc	4.8 a-c	0.8 ab	3.0 a	92.0 d-g	0.0 b	3.3 e-j
N17045	5.6 ab	4.8 a-c	0.9 a	2.2 b-g	92.6 c-e	0.0 ab	3.5 e-j
N17047	5.6 a-c	4.8 ab	0.8 ab	2.3 b-f	92.8 b-d	0.1 ab	3.1 h-j
Mean	5.6	4.8	0.8	2.3	92.1	0.1	3.8
LSD¹	0.1	0.1	0.1	0.7	1.2	0.1	1.0

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 9. Laboratory sample blanching of Extra-Large Kernels (ELK). Averages from Tidewater AREC (Suffolk, VA) and Martin County, NC. Two-year averages (2019- 2020).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 a	4.8 a	0.9 a	3.0 a-d	92.5 a-c	0.0 ab	3.0 ef
Bailey II	5.7 a	4.8 a	0.9 a	2.0 cd	93.1 ab	0.0 ab	3.3 d-f
Emery	5.7 a	4.7 a	0.9 a	1.6 d	94.4 a	0.0 ab	2.5 f
Sullivan	5.7 a	4.8 a	0.9 a	2.9 a-d	91.6 b-d	0.0 b	3.9 b-f
Walton	5.7 a	4.7 a	0.9 a	4.2 a	88.8 e	0.0 ab	5.4 a-c
Wynne	5.7 a	4.8 a	0.9 a	2.3 b-d	91.2 b-d	0.1 ab	4.8 a-d
N14002oIJ	5.6 a	4.8 a	0.9 a	3.2 a-c	90.4 c-e	0.0 b	4.8 a-d
N14004oIJ	5.7 a	4.7 a	0.9 a	3.4 ab	90.4 c-e	0.0 ab	4.6 a-c
N14023oI	5.7 a	4.8 a	0.9 a	3.4 ab	89.5 de	0.1 a	5.5 ab
N14027oIJ	5.7 a	4.8 a	0.9 a	2.8 a-d	89.4 de	0.0 ab	6.1 a
N15017oI	5.7 a	4.7 a	0.9 a	3.1 a-c	92.1 bc	0.0 ab	3.3 d-f
N15039oI	5.7 a	4.7 a	0.9 a	3.6 ab	91.2 b-d	0.0 b	3.7 c-f
N15041oI	5.7 a	4.7 a	0.9 a	3.3 a-c	89.5 de	0.1 ab	5.6 ab
N15044oIF	5.7 a	4.7 a	0.9 a	3.1 a-c	89.4 de	0.1 ab	5.9 a
N16005	5.7 a	4.8 a	0.9 a	3.1 a-c	92.0 bc	0.0 ab	3.4 d-f
Mean	5.7	4.7	0.9	3.0	91.0	0.0	4.4
LSD¹	0.1	0.1	0.1	1.4	2.3	0.1	1.7

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 10. Laboratory sample blanching of Extra-Large Kernels (ELK). Averages from Tidewater AREC (Suffolk, VA) and Martin County, NC. Three-year averages (2018- 2020).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 a	4.8 a	0.9 a	2.4 b-e	93.0 ab	0.0 ab	3.0 de
Bailey II	5.7 a	4.8 a	0.9 a	2.0 de	92.9 bc	0.0 ab	3.4 c-e
Emery	5.7 a	4.8 a	0.9 a	1.5 e	94.6 a	0.0 ab	2.3 e
Sullivan	5.7 a	4.8 a	0.9 a	2.5 b-e	92.3 bc	0.0 b	3.6 c-e
Walton	5.7 a	4.8 a	0.9 a	4.1 a	88.9 g	0.0 ab	5.4 ab
Wynne	5.7 a	4.8 a	0.9 a	2.2 c-e	91.8 b-d	0.0 ab	4.4 a-c
N14002oIJ	5.7 a	4.8 a	0.9 a	2.8 b-d	91.5 b-e	0.0 b	4.2 b-d
N14004oIJ	5.7 a	4.8 a	0.9 a	3.0 b-d	91.2 c-f	0.0 ab	4.2 b-d
N14023oI	5.7 a	4.8 a	0.9 a	3.1 a-c	90.2 d-g	0.1 a	5.0 ab
N14027oIJ	5.7 a	4.8 a	0.9 a	2.8 b-d	90.1 e-g	0.0 ab	5.5 a
N15017oI	5.7 a	4.8 a	0.9 a	2.8 b-d	92.5 bc	0.0 ab	3.1 de
N15039oI	5.7 a	4.8 a	0.9 a	3.3 ab	91.4 b-e	0.0 b	3.7 cd
N15041oI	5.7 a	4.8 a	0.9 a	3.3 ab	89.7 fg	0.0 ab	5.4 ab
N15044oIF	5.7 a	4.8 a	0.9 a	3.0 b-d	90.2 d-g	0.0 ab	5.1 ab
Mean	5.7	4.8	0.9	2.8	91.4	0.0	4.2
LSD¹	0.1	0.1	0.1	1.0	1.7	0.1	1.3

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 11. Laboratory sample blanching of Medium Kernels from Tidewater AREC (Suffolk, VA), Dig 1, 2020 (28 September).

Variety	% H ₂ O before roasting	% H ₂ O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.8 a	4.9 ab	0.9 a	3.0 a-d	86.0 ab	1.5 cd	8.0 h
Bailey II	5.8 a	4.9 ab	0.9 a	2.0 d	87.1 a	1.3 d	8.0 h
Emery	5.8 a	4.9 ab	0.9 a	2.3 b-d	84.6 a-c	2.7 ab	8.8 gh
Sullivan	5.7 a	4.9 ab	0.8 a	2.1 cd	84.1 bc	2.3 a-c	10.0 f-h
Walton	5.8 a	4.9 ab	0.9 a	3.1 a-d	80.4 d-j	1.8 b-d	13.2 a-d
Wynne	5.7 a	4.9 ab	0.8 a	2.4 b-d	81.8 c-h	2.3 a-c	11.9 c-f
N14001	5.7 a	5.0 a	0.7 a	3.3 a-d	82.7 c-e	2.0 a-d	10.4 e-h
N14002oIJ	5.8 a	5.0 a	0.8 a	3.2 a-d	80.2 d-k	2.1 a-d	12.9 a-e
N14004oIJ	5.8 a	4.9 ab	0.9 a	2.4 b-d	82.6 c-f	2.6 ab	10.9 d-g
N14007	5.8 a	5.0 ab	0.8 a	3.3 a-d	79.2 h-k	2.3 a-c	13.5 a-c
N14009	5.8 a	5.0 ab	0.8 a	3.1 a-d	82.7 c-e	2.3 a-c	10.4 e-h
N15066	5.7 a	4.9 ab	0.8 a	3.0 a-d	80.1 e-k	2.9 a	12.5 a-f
N14023oI	5.8 a	5.0 a	0.8 a	3.0 a-d	78.6 i-k	2.2 a-c	14.6 ab
N14027oIJ	5.8 a	5.0 ab	0.8 a	3.1 a-d	81.1 d-i	2.2 a-c	12.0 b-f
N15017oI	5.7 a	4.9 ab	0.8 a	3.3 a-d	77.4 k	2.9 a	14.9 a
N15039oI	5.8 a	5.0 ab	0.8 a	3.2 a-d	79.4 h-k	2.5 ab	13.3 a-d
N15041oI	5.7 a	4.9 b	0.9 a	3.1 a-d	78.7 i-k	2.7 a	13.9 a-c
N15044oIF	5.8 a	4.9 ab	0.9 a	3.5 ab	78.7 i-k	2.5 ab	13.7 a-c
N15053	5.7 a	4.9 ab	0.8 a	3.8 a	78.7 i-k	2.0 a-d	13.9 a-c
N15060	5.7 a	4.9 ab	0.8 a	3.4 a-c	80.9 d-i	2.3 a-c	11.8 c-f
N16005	5.8 a	4.9 ab	0.9 a	3.4 ab	79.7 f-k	2.5 ab	13.3 a-d
N16012	5.8 a	4.9 ab	0.9 a	3.8 a	78.3 i-k	2.4 a-c	13.9 a-c
N16021	5.8 a	5.0 ab	0.8 a	3.9 a	79.2 h-k	2.3 a-c	13.1 a-d
N17036	5.7 a	4.9 b	0.8 a	3.5 ab	78.3 i-k	2.5 ab	14.1 a-c
N17037	5.7 a	4.9 ab	0.8 a	2.7 a-d	83.0 cd	2.3 a-c	10.4 e-h
N17040	5.7 a	4.9 ab	0.8 a	3.5 ab	77.9 jk	2.6 ab	14.5 ab
N17041	5.7 a	4.9 b	0.8 a	2.7 a-d	79.3 h-k	2.9 a	13.5 a-c
N17044	5.7 a	5.0 ab	0.8 a	2.8 a-d	79.6 g-k	2.5 ab	13.5 a-c
N17045	5.8 a	5.0 a	0.8 a	2.6 a-d	82.4 c-g	2.5 ab	10.9 d-g
N17047	5.7 a	4.9 ab	0.8 a	3.6 ab	80.2 d-k	2.1 a-d	12.6 a-f
Mean	5.7	4.9	0.8	3.0	80.7	2.3	12.3
LSD¹	0.1	0.1	0.2	1.3	2.3	0.9	2.6

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 12. Laboratory sample blanching of Medium Kernels from Tidewater AREC (Suffolk, VA), Dig 2, 2020 (20 October).

Variety	% H ₂ O before roasting	% H ₂ O after roasting	%Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 a	4.9 a-c	0.8 ab	3.0 b-d	82.5 a	2.5 e	10.3 i
Bailey II	5.6 a	4.8 bc	0.8 ab	2.5 cd	81.1 ab	2.8 c-e	12.0 f-i
Emery	5.7 a	4.8 bc	0.9 ab	3.2 a-d	81.2 ab	3.5 a-e	10.5 hi
Sullivan	5.7 a	4.9 a-c	0.8 ab	3.0 b-d	79.2 a-d	3.2 b-e	13.0 b-i
Walton	5.6 a	4.8 bc	0.8 ab	3.0 b-d	80.0 ab	3.3 a-e	12.1 e-i
Wynne	5.7 a	4.9 a-c	0.8 ab	3.3 a-d	78.2 b-e	3.9 ab	13.0 b-i
N14001	5.7 a	4.9 a-c	0.8 ab	2.5 cd	80.3 ab	2.9 b-e	12.8 c-i
N14002oIJ	5.7 a	4.9 a-c	0.8 ab	3.5 a-d	79.5 a-c	3.2 b-e	12.2 e-i
N14004oIJ	5.7 a	4.9 a-c	0.9 ab	2.5 cd	81.0 ab	2.5 e	12.5 d-i
N14007	5.7 a	4.9 a-c	0.9 ab	3.5 a-d	77.9 b-e	2.9 b-e	14.2 a-g
N14009	5.7 a	4.8 bc	0.9 ab	4.1 ab	73.0 f	4.3 a	17.0 a
N15066	5.6 a	4.8 c	0.9 ab	3.7 a-c	75.1 ef	3.7 a-c	15.9 a-c
N14023oI	5.6 a	4.9 a-c	0.8 b	3.9 ab	75.0 ef	3.3 a-e	16.2 ab
N14027oIJ	5.7 a	4.9 ab	0.8 b	4.2 ab	75.1 d-f	3.5 a-e	15.6 a-d
N15017oI	5.7 a	4.8 c	0.9 a	4.0 ab	75.7 c-f	3.4 a-e	15.4 a-e
N15039oI	5.6 a	4.8 bc	0.8 ab	3.7 a-c	77.1 b-e	3.4 a-e	14.1 a-g
N15041oI	5.7 a	4.8 c	0.9 a	4.4 a	75.2 d-f	4.0 ab	14.9 a-f
N15044oIF	5.6 a	4.8 c	0.9 ab	3.6 a-d	78.0 b-e	2.9 b-e	13.9 a-g
N15053	5.7 a	4.8 bc	0.9 ab	4.1 ab	75.7 c-f	3.7 a-d	14.9 a-f
N15060	5.6 a	4.8 bc	0.8 ab	3.4 a-d	75.7 c-f	3.7 a-c	15.6 a-d
N16005	5.7 a	4.8 bc	0.9 ab	3.5 a-d	79.0 a-e	2.6 de	13.4 b-i
N16012	5.7 a	4.8 bc	0.9 ab	4.2 ab	77.5 b-e	3.3 a-e	13.5 b-i
N16021	5.6 a	4.8 bc	0.8 ab	3.6 a-d	77.4 b-e	3.3 a-e	14.0 a-g
N17036	5.6 a	4.9 a-c	0.8 b	3.4 a-d	78.3 b-e	2.9 b-e	13.8 a-h
N17037	5.6 a	4.8 bc	0.8 ab	2.9 b-d	79.1 a-e	2.9 b-e	13.6 b-i
N17040	5.6 a	4.8 bc	0.8 ab	3.0 b-d	79.7 a-c	2.9 b-e	12.8 c-i
N17041	5.6 a	4.8 bc	0.8 ab	3.5 a-d	78.1 b-e	3.9 a-c	13.0 b-i
N17044	5.7 a	5.0 a	0.8 b	3.1 a-d	78.0 b-e	3.2 b-e	14.0 a-g
N17045	5.7 a	4.9 a-c	0.8 ab	3.2 a-d	81.0 ab	3.8 a-c	10.9 g-i
N17047	5.7 a	4.8 bc	0.9 ab	2.4 d	78.7 a-e	3.6 a-e	13.8 a-h
Mean	5.6	4.8	0.8	3.4	89.8	3.3	13.6
LSD¹	0.1	0.1	0.1	1.3	4.1	1.1	3.3

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 13. Laboratory sample blanching of Medium Kernels. Averages from both digging dates from Tidewater AREC (Suffolk, VA), 2020.

Variety	% H ₂ O before roasting	% H ₂ O after roasting	%Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.7 a	4.9 a-c	0.8 a-c	3.0 c-g	84.2 a	2.0 b	9.1 g
Bailey II	5.7 a	4.9 a-c	0.8 a-c	2.3 g	84.1 a	2.0 b	10.0 e-g
Emery	5.7 a	4.9 a-c	0.9 ab	2.8 d-g	82.9 ab	3.1 ab	9.7 fg
Sullivan	5.7 a	4.9 a-c	0.8 a-c	2.5 e-g	81.6 a-d	2.7 ab	11.5 c-g
Walton	5.7 a	4.9 a-c	0.8 a-c	3.0 b-g	80.2 b-g	2.5 ab	12.6 b-d
Wynne	5.7 a	4.9 a-c	0.8 a-c	2.8 c-g	80.0 b-g	3.1 a	12.4 b-e
N14001	5.7 a	4.9 ab	0.8 c	2.9 c-g	81.5 a-e	2.4 ab	11.6 c-g
N14002oIJ	5.7 a	4.9 ab	0.8 bc	3.3 a-f	79.9 b-h	2.6 ab	12.5 b-d
N14004oIJ	5.7 a	4.9 a-c	0.9 ab	2.4 fg	81.8 a-c	2.6 ab	11.7 c-f
N14007	5.7 a	4.9 a-c	0.8 a-c	3.4 a-e	78.5 c-h	2.6 ab	13.8 a-c
N14009	5.7 a	4.9 a-c	0.8 a-c	3.6 a-d	77.8 f-h	3.3 a	13.7 a-c
N15066	5.7 a	4.8 bc	0.8 a-c	3.3 a-e	77.6 gh	3.3 a	14.2 ab
N14023oI	5.7 a	4.9 ab	0.8 c	3.4 a-d	76.8 gh	2.8 ab	15.4 a
N14027oIJ	5.7 a	4.9 ab	0.8 bc	3.6 a-d	78.1 e-h	2.9 ab	13.8 a-c
N15017oI	5.7 a	4.8 bc	0.9 ab	3.6 a-d	76.5 h	3.1 a	15.1 a
N15039oI	5.7 a	4.9 a-c	0.8 a-c	3.5 a-d	78.2 d-h	3.0 ab	13.7 a-c
N15041oI	5.7 a	4.8 c	0.9 a	3.7 a-c	76.9 gh	3.3 a	14.4 ab
N15044oIF	5.7 a	4.8 bc	0.9 ab	3.5 a-d	78.4 c-h	2.7 ab	13.8 a-c
N15053	5.7 a	4.9 a-c	0.8 a-c	3.9 ab	77.2 gh	2.8 ab	14.4 ab
N15060	5.7 a	4.9 a-c	0.8 a-c	3.4 a-e	78.3 d-h	3.0 ab	13.7 a-c
N16005	5.7 a	4.9 a-c	0.9 ab	3.4 a-d	79.4 c-h	2.5 ab	13.4 a-c
N16012	5.7 a	4.9 a-c	0.9 ab	4.0 a	77.9 f-h	2.9 ab	13.7 a-c
N16021	5.7 a	4.9 a-c	0.8 a-c	3.7 a-c	78.3 d-h	2.8 ab	13.5 a-c
N17036	5.6 a	4.9 a-c	0.8 bc	3.4 a-d	78.3 d-h	2.7 ab	13.9 a-c
N17037	5.7 a	4.9 a-c	0.8 a-c	2.8 d-g	81.0 a-f	2.6 ab	12.0 b-f
N17040	5.7 a	4.9 a-c	0.8 a-c	3.2 a-f	78.8 c-h	2.7 ab	13.7 a-c
N17041	5.6 a	4.8 bc	0.8 a-c	3.1 a-g	78.7 c-h	3.4 a	13.2 a-d
N17044	5.7 a	5.0 a	0.8 c	2.9 c-g	78.8 c-h	2.8 ab	13.8 a-c
N17045	5.7 a	4.9 ab	0.8 bc	2.9 c-g	81.7 a-d	3.2 a	10.9 d-g
N17047	5.7 a	4.9 a-c	0.8 a-c	3.0 c-g	79.4 c-h	2.8 ab	13.2 a-d
Mean	5.7	4.9	0.8	3.2	79.4	2.8	12.9
LSD¹	0.1	0.1	0.1	0.9	3.4	1.1	2.5

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 14. Laboratory sample blanching of Medium Kernels from Martin County, NC, Dig 1, 2020 (5 October).

Variety	% H ₂ O before roasting	% H ₂ O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.9 a	5.0 ab	0.9 a	3.0 c-g	83.8 a	1.8 fg	9.8 ij
Bailey II	5.9 a	5.0 ab	0.9 a	3.0 c-g	84.3 a	1.8 fg	9.4 j
Emery	5.8 a	4.9 ab	0.9 a	2.3 g	82.6 a-c	2.0 c-g	11.6 e-j
Sullivan	5.8 a	4.9 ab	0.9 a	3.2 b-g	82.3 a-f	1.9 e-g	11.1 f-j
Walton	5.8 a	4.9 ab	0.9 a	3.8 a-e	80.7 b-i	2.3 b-g	11.6 e-j
Wynne	5.8 a	5.0 ab	0.9 a	3.1 b-g	79.6 f-k	2.7 a-g	13.1 c-h
N14001	5.8 a	4.9 ab	0.9 a	3.4 a-f	82.5 a-d	1.8 g	10.8 h-j
N14002oIJ	5.9 a	5.0 ab	0.9 a	3.3 a-f	80.4 c-j	2.9 a-d	11.9 d-i
N14004oIJ	5.8 a	4.9 ab	0.9 a	2.7 fg	78.8 i-m	2.8 a-e	14.1 a-d
N14007	5.8 a	5.0 ab	0.9 a	4.1 ab	77.9 j-n	2.3 b-g	14.1 a-e
N14009	5.8 a	5.0 ab	0.9 a	3.8 a-e	78.3 i-n	2.9 a-d	13.5 b-f
N15066	5.8 a	4.9 ab	0.9 a	3.2 b-g	77.6 k-n	3.6 a	14.1 a-e
N14023oI	5.8 a	4.9 ab	0.9 a	4.3 a	77.2 k-n	2.9 a-d	14.1 a-d
N14027oIJ	5.8 a	4.9 b	0.9 a	3.0 d-g	78.7 i-m	2.6 b-g	14.2 a-d
N15017oI	5.8 a	4.9 ab	0.9 a	3.5 a-f	79.2 h-l	2.4 b-g	13.3 b-g
N15039oI	5.9 a	5.0 ab	0.9 a	2.8 e-g	82.4 a-e	2.4 b-g	10.9 g-j
N15041oI	5.8 a	5.0 ab	0.9 a	3.6 a-f	75.8 n	2.7 a-g	16.3 a
N15044oIF	5.8 a	4.9 ab	0.9 a	3.2 b-g	78.6 i-m	2.8 a-e	13.8 b-e
N15053	5.8 a	4.9 ab	0.9 a	3.8 a-e	78.1 i-n	2.8 a-f	13.8 b-e
N15060	5.8 a	5.0 ab	0.9 a	3.5 a-f	78.1 i-n	2.9 a-c	13.7 b-e
N16005	5.8 a	4.9 ab	0.9 a	3.5 a-f	82.0 a-g	1.9 e-g	11.1 f-j
N16012	5.8 a	4.9 ab	0.9 a	3.9 a-d	79.7 e-k	2.6 b-g	12.2 d-i
N16021	5.8 a	4.9 ab	0.9 a	3.0 c-g	79.5 g-k	2.8 a-e	13.2 c-h
N17036	5.8 a	5.0 ab	0.9 a	3.1 b-g	79.8 d-k	2.5 b-g	13.0 c-h
N17037	5.9 a	5.0 a	0.9 a	3.6 a-f	81.9 a-h	2.5 b-g	10.4 ij
N17040	5.8 a	4.9 ab	0.9 a	3.2 b-g	76.5 mn	3.1 ab	15.7 ab
N17041	5.8 a	5.0 a	0.8 a	3.0 d-g	78.6 i-m	2.8 a-f	14.1 a-e
N17044	5.8 a	4.9 b	0.9 a	4.0 a-c	76.7 l-n	3.0 ab	14.8 a-c
N17045	5.8 a	5.0 ab	0.9 a	3.2 b-g	83.2 ab	1.9 d-g	10.1 ij
N17047	5.8 a	4.9 ab	0.9 a	3.6 a-f	78.2 i-n	3.0 ab	13.6 b-e
Mean	5.8	4.9	0.9	3.3	79.8	2.5	12.8
LSD¹	0.1	0.1	0.1	1.0	2.7	1.0	2.5

¹Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 15. Laboratory sample blanching of Medium Kernels from Martin County, NC, Dig 2, 2020 (15 October).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.8 a	4.9 ab	0.9 ab	3.7 a-c	81.8 a	2.1 f	10.9 j
Bailey II	5.8 a	4.8 b	1.0 a	3.4 a-c	81.2 a-c	2.3 ef	11.5 ij
Emery	5.8 a	4.9 ab	0.9 ab	3.7 a-c	81.4 ab	2.3 ef	10.8 j
Sullivan	5.8 a	4.9 ab	0.9 ab	3.3 a-c	78.5 b-i	2.8 b-f	13.9 b-h
Walton	5.8 a	4.9 ab	0.9 ab	3.3 a-c	78.2 d-k	2.8 b-f	14.2 b-g
Wynne	5.8 a	4.9 ab	0.9 ab	3.2 a-c	78.9 a-h	2.6 c-f	13.8 b-h
N14001	5.8 a	4.9 ab	0.9 ab	3.3 a-c	79.6 a-f	2.8 b-f	12.7 f-j
N14002oIJ	5.8 a	4.9 ab	0.9 ab	3.0 a-c	77.2 f-m	3.9 ab	14.4 a-g
N14004oIJ	5.9 a	5.0 a	0.9 ab	3.2 a-c	76.9 f-m	3.3 a-f	15.0 a-f
N14007	5.8 a	4.9 ab	0.9 ab	3.7 a-c	77.3 f-m	3.2 a-f	14.3 b-g
N14009	5.8 a	4.9 ab	0.9 ab	3.9 ab	78.0 e-l	3.0 b-f	13.5 c-i
N15066	5.8 a	4.9 ab	0.9 ab	2.9 a-c	78.6 b-i	3.6 a-e	13.4 d-i
N14023oI	5.8 a	5.0 ab	0.9 ab	3.7 a-c	78.8 b-h	2.1 f	13.9 b-h
N14027oIJ	5.8 a	4.9 ab	0.9 ab	2.8 bc	78.4 c-j	3.0 b-f	14.3 a-g
N15017oI	5.8 a	4.9 ab	0.9 ab	3.5 a-c	76.3 h-n	3.4 a-e	15.2 a-e
N15039oI	5.8 a	4.9 ab	0.9 ab	3.1 a-c	80.6 a-e	2.4 d-f	12.4 g-j
N15041oI	5.8 a	4.9 ab	0.9 ab	4.2 a	73.8 n	3.9 a-c	16.6 a
N15044oIF	5.7 a	4.9 ab	0.9 ab	3.5 a-c	75.1 l-n	4.0 ab	15.9 ab
N15053	5.8 a	4.9 ab	0.9 ab	4.0 ab	76.4 g-n	3.0 b-f	14.9 a-f
N15060	5.7 a	4.8 b	0.9 ab	3.8 ab	75.6 i-n	3.1 b-f	15.9 ab
N16005	5.8 a	5.0 ab	0.8 b	3.7 a-c	81.0 a-d	2.4 d-f	11.3 ij
N16012	5.7 a	4.9 ab	0.9 ab	4.0 ab	76.0 h-n	3.0 b-f	15.4 a-d
N16021	5.7 a	4.9 ab	0.8 b	4.2 a	75.3 k-n	3.6 a-e	15.3 a-d
N17036	5.8 a	5.0 ab	0.8 b	3.9 ab	75.4 j-n	3.4 a-f	15.8 a-c
N17037	5.8 a	4.9 ab	0.9 ab	3.7 a-c	78.6 b-i	2.9 b-f	13.3 d-i
N17040	5.7 a	4.9 ab	0.8 b	3.5 a-c	74.6 mn	4.5 a	15.8 a-c
N17041	5.8 a	4.9 ab	0.9 ab	3.5 a-c	76.7 f-n	3.8 a-c	14.5 a-g
N17044	5.7 a	4.9 ab	0.9 ab	2.4 c	80.4 a-e	3.7 a-d	11.9 h-j
N17045	5.8 a	4.9 ab	0.9 ab	3.2 a-c	78.1 d-k	3.8 a-c	13.3 d-i
N17047	5.8 a	4.9 ab	0.9 ab	3.3 a-c	79.4 a-g	2.7 b-f	13.0 e-j
Mean	5.7	4.9	0.9	3.5	77.9	3.1	13.9
LSD²	0.2	0.2	0.1	1.3	3.0	1.3	2.3

¹Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 16. Laboratory sample blanching of Medium Kernels. Averages from both digging dates from Martin County, NC, 2020.

Variety	% H ₂ O before roasting	% H ₂ O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.8 a	4.9 ab	0.9 ab	3.4 a-c	82.8 a	1.9 i	10.3 i
Bailey II	5.8 ab	4.9 ab	0.9 a	3.2 a-c	82.7 a	2.1 hi	10.4 i
Emery	5.8 ab	4.9 ab	0.9 ab	3.0 c	82.0 ab	2.1 g-i	11.2 hi
Sullivan	5.8 ab	4.9 ab	0.9 a-c	3.2 a-c	80.4 a-g	2.3 e-i	12.5 d-h
Walton	5.8 ab	4.9 ab	0.9 ab	3.6 a-c	79.4 b-i	2.5 c-i	12.9 c-h
Wynne	5.8 ab	4.9 ab	0.9 a-c	3.1 bc	79.3 c-i	2.6 c-i	13.4 c-g
N14001	5.8 ab	4.9 ab	0.9 a-c	3.3 a-c	81.1 a-e	2.3 f-i	11.8 f-i
N14002oIJ	5.8 ab	4.9 ab	0.9 a-c	3.1 bc	78.8 d-i	3.4 a-c	13.1 c-h
N14004oIJ	5.8 ab	5.0 a	0.9 bc	3.0 c	77.8 g-j	3.1 a-f	14.5 a-c
N14007	5.8 ab	4.9 ab	0.9 bc	3.9 ab	77.6 h-j	2.8 b-i	14.2 b-d
N14009	5.8 ab	4.9 ab	0.9 bc	3.9 ab	78.1 f-j	2.9 a-h	13.5 c-g
N15066	5.8 ab	4.9 ab	0.9 bc	3.0 c	78.1 f-j	3.6 ab	13.7 b-f
N14023oI	5.8 ab	4.9 ab	0.9 a-c	4.0 a	78.0 f-j	2.5 d-i	14.0 b-d
N14027oIJ	5.8 ab	4.9 ab	0.9 a-c	2.9 c	78.5 e-i	2.8 b-i	14.2 b-d
N15017oI	5.8 ab	4.9 ab	0.9 bc	3.5 a-c	77.7 g-j	2.9 a-h	14.3 b-d
N15039oI	5.8 ab	4.9 ab	0.9 a-c	2.9 c	81.5 a-d	2.4 d-i	11.6 g-i
N15041oI	5.8 ab	4.9 ab	0.9 bc	3.9 ab	74.8 k	3.3 a-d	16.4 a
N15044oIF	5.8 ab	4.9 ab	0.9 a-c	3.4 a-c	76.8 i-k	3.4 a-c	14.8 a-c
N15053	5.8 ab	4.9 ab	0.9 ab	3.9 ab	77.3 i-k	2.9 b-h	14.4 b-d
N15060	5.8 ab	4.9 ab	0.9 a-c	3.7 a-c	76.9 i-k	3.0 a-g	14.8 a-c
N16005	5.8 ab	4.9 ab	0.9 bc	3.6 a-c	81.5 a-c	2.1 g-i	11.2 hi
N16012	5.8 ab	4.9 ab	0.9 a-c	4.0 a	77.8 g-j	2.8 b-i	13.8 b-e
N16021	5.7 b	4.9 ab	0.8 c	3.6 a-c	77.4 i-k	3.2 a-e	14.2 b-d
N17036	5.8 ab	5.0 a	0.8 c	3.5 a-c	77.6 h-j	2.9 a-h	14.4 b-d
N17037	5.8 ab	4.9 ab	0.9 a-c	3.7 a-c	80.3 a-h	2.7 c-i	11.8 e-i
N17040	5.7 b	4.9 ab	0.8 c	3.4 a-c	75.5 jk	3.8 a	15.7 ab
N17041	5.8 ab	5.0 a	0.8 c	3.2 a-c	77.6 h-j	3.3 a-d	14.3 b-d
N17044	5.7 b	4.9 b	0.9 a-c	3.2 a-c	78.5 e-i	3.3 a-d	13.4 c-g
N17045	5.8 ab	4.9 ab	0.9 a-c	3.2 a-c	80.7 a-f	2.9 b-h	11.7 g-i
N17047	5.8 ab	4.9 ab	0.9 bc	3.4 a-c	78.8 d-i	2.8 b-i	13.3 c-g
Mean	5.8	4.9	0.9	3.4	78.8	2.8	13.3
LSD¹	0.1	0.1	0.1	0.8	2.7	0.9	2.0

¹Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 17. Laboratory sample blanching of Medium Kernels. Averages from Tidewater AREC (Suffolk, VA) and Martin County, NC, 2020.

Variety	% H ₂ O before roasting	% H ₂ O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.8 a	4.9 a	0.9 ab	3.2 c-j	83.5 a	2.0 h	9.7 l
Bailey II	5.7 a	4.9 a	0.9 a	2.7 ij	83.4 a	2.0 gh	10.2 kl
Emery	5.8 a	4.9 a	0.9 a	2.9 h-j	82.5 ab	2.6 c-h	10.4 j-l
Sullivan	5.7 a	4.9 a	0.8 ab	2.9 h-j	81.0 bc	2.5 d-h	12.0 g-j
Walton	5.7 a	4.9 a	0.9 ab	3.3 b-i	79.8 c-f	2.5 d-h	12.8 d-i
Wynne	5.7 a	4.9 a	0.8 ab	3.0 g-j	79.6 c-g	2.8 a-f	12.9 c-h
N14001	5.7 a	4.9 a	0.8 ab	3.1 d-j	81.3 bc	2.3 e-h	11.7 h-k
N14002oIJ	5.8 a	4.9 a	0.8 ab	3.2 b-j	79.3 c-h	3.0 a-e	12.8 c-i
N14004oIJ	5.8 a	4.9 a	0.9 ab	2.7 j	79.8 c-f	2.8 a-f	13.1 c-h
N14007	5.8 a	4.9 a	0.8 ab	3.6 a-f	78.1 f-i	2.7 b-g	14.0 a-e
N14009	5.7 a	4.9 a	0.8 ab	3.7 a-c	78.0 f-j	3.1 a-d	13.6 b-g
N15066	5.7 a	4.9 a	0.8 ab	3.2 c-j	77.8 f-j	3.4 a	14.0 a-e
N14023oI	5.7 a	4.9 a	0.8 ab	3.7 a-d	77.4 h-j	2.6 c-h	14.7 ab
N14027oIJ	5.7 a	4.9 a	0.8 ab	3.2 b-j	78.3 e-i	2.8 a-f	14.0 a-e
N15017oI	5.7 a	4.9 a	0.9 ab	3.6 a-g	77.1 ij	3.0 a-f	14.7 ab
N15039oI	5.7 a	4.9 a	0.8 ab	3.2 b-j	79.8 c-f	2.7 b-g	12.7 e-i
N15041oI	5.7 a	4.9 a	0.9 ab	3.8 ab	75.9 j	3.3 ab	15.4 a
N15044oIF	5.7 a	4.9 a	0.9 ab	3.4 a-h	77.6 g-j	3.0 a-d	14.3 a-d
N15053	5.7 a	4.9 a	0.9 ab	3.9 a	77.2 h-j	2.8 a-f	14.4 a-c
N15060	5.7 a	4.9 a	0.8 ab	3.5 a-g	77.6 g-j	3.0 a-e	14.2 a-d
N16005	5.7 a	4.9 a	0.9 ab	3.5 a-g	80.4 b-e	2.3 f-h	12.3 f-i
N16012	5.7 a	4.9 a	0.9 ab	4.0 a	77.9 f-j	2.8 a-f	13.7 b-f
N16021	5.7 a	4.9 a	0.8 ab	3.7 a-e	77.8 f-j	3.0 a-e	13.9 a-e
N17036	5.7 a	4.9 a	0.8 b	3.5 a-h	77.9 f-j	2.8 a-f	14.1 a-e
N17037	5.7 a	4.9 a	0.8 ab	3.2 b-j	80.6 b-d	2.6 c-h	11.9 h-j
N17040	5.7 a	4.9 a	0.8 ab	3.3 b-j	77.1 ij	3.3 a-c	14.7 ab
N17041	5.7 a	4.9 a	0.8 ab	3.2 c-j	78.1 f-i	3.3 ab	13.7 b-f
N17044	5.7 a	4.9 a	0.8 ab	3.1 e-j	78.7 d-i	3.1 a-d	13.6 b-g
N17045	5.7 a	4.9 a	0.8 ab	3.0 f-j	81.2 bc	3.0 a-e	11.3 i-l
N17047	5.7 a	4.9 a	0.8 ab	3.2 c-j	79.1 c-i	2.8 a-f	13.2 b-h
Mean	5.7	4.9	0.8	3.3	79.1	2.8	13.1
LSD¹	0.1	0.1	0.1	0.6	2.2	0.7	1.6

¹Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 18. Laboratory sample blanching of Medium Kernels. Averages from Tidewater AREC (Suffolk, VA) and Martin County, NC. Two-year averages (2019- 2020).

Variety	% H ₂ O before roasting	% H ₂ O after roasting	%Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.8 a	4.9 a	1.0 a	2.9 b-e	84.4 ab	1.7 cd	9.3 i
Bailey II	5.8 a	4.9 a	1.0 a	2.6 e	84.8 a	1.7 d	9.4 hi
Emery	5.8 a	4.9 a	0.9 a	2.6 e	82.4 b-d	2.4 ab	10.9 f-h
Sullivan	5.8 a	4.9 a	0.9 a	2.8 de	82.8 a-c	2.1 b-d	10.5 g-i
Walton	5.8 a	4.9 a	0.9 a	3.7 a	79.6 e-h	2.6 ab	12.5 b-e
Wynne	5.8 a	4.9 a	0.9 a	2.9 c-e	81.6 c-e	2.3 a-c	11.6 e-g
N14002oIJ	5.8 a	4.9 a	0.9 a	3.3 a-d	80.6 d-f	2.6 ab	11.9 d-g
N14004oIJ	5.8 a	4.9 a	0.9 a	2.8 de	80.3 e-g	2.3 ab	13.0 a-e
N14023oI	5.8 a	4.9 a	0.9 a	3.6 a	78.4 gh	2.5 ab	13.9 a-c
N14027oIJ	5.8 a	4.9 a	0.9 a	3.5 ab	78.6 gh	2.3 ab	14.0 ab
N15017oI	5.8 a	4.9 a	0.9 a	3.5 a	78.9 f-h	2.6 ab	13.3 a-d
N15039oI	5.8 a	4.9 a	0.9 a	3.4 ab	80.8 c-f	2.5 ab	11.7 e-g
N15041oI	5.8 a	4.9 a	0.9 a	3.4 a-c	78.1 h	2.7 a	14.2 a
N15044oIF	5.8 a	4.9 a	0.9 a	3.6 a	78.1 h	2.6 ab	14.1 ab
N16005	5.8 a	4.9 a	0.9 a	3.8 a	79.8 e-h	2.4 ab	12.4 c-f
Mean	5.8	4.9	0.9	3.2	80.6	2.3	12.2
LSD¹	0.1	0.1	0.1	0.5	2.0	0.5	1.6

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Blanching Results

Table 19. Laboratory sample blanching of Medium Kernels. Averages from Tidewater AREC (Suffolk, VA) and Martin County, NC. Three-year averages (2018 - 2020).

Variety	% H2O before roasting	% H2O after roasting	% Blanching loss	% Splits Blanched	% Whole Blanched	% Not Blanched	% Partially Blanched
Bailey	5.8 a	4.9 a	0.9 ab	3.1 ef	84.3 a	1.5 e	9.4 f
Bailey II	5.8 a	4.8 a	0.9 a	2.9 ef	84.1 ab	1.7 de	9.7 f
Emery	5.8 a	4.9 a	0.9 a-c	2.8 f	82.8 bc	2.2 a-d	10.5 ef
Sullivan	5.8 a	4.9 a	0.9 a-c	3.0 ef	83.0 a-c	1.8 c-e	10.5 ef
Walton	5.8 a	4.9 a	0.9 a-c	4.3 a	79.3 d-f	2.5 a	12.3 bc
Wynne	5.8 a	4.8 a	0.9 a-c	2.9 ef	82.5 c	1.9 c-e	11.1 de
N14002oIJ	5.8 a	4.9 a	0.9 a-c	3.4 c-e	80.7 d	2.4 ab	11.9 cd
N14004oIJ	5.8 a	4.9 a	0.9 a-c	3.2 d-f	80.2 de	2.0 b-d	12.9 a-c
N14023oI	5.7 a	4.9 a	0.9 c	3.7 b-d	78.6 f	2.3 ab	13.7 a
N14027oIJ	5.7 a	4.8 a	0.9 a-c	3.7 b-d	78.8 ef	2.2 a-c	13.7 a
N15017oI	5.7 a	4.8 a	0.9 a-c	3.9 a-c	78.6 f	2.5 a	13.3 ab
N15039oI	5.7 a	4.9 a	0.9 bc	3.8 a-c	80.3 de	2.3 ab	11.9 cd
N15041oI	5.8 a	4.8 a	0.9 ab	3.8 a-c	78.1 f	2.5 a	13.9 a
N15044oIF	5.7 a	4.8 a	0.9 a-c	4.1 ab	78.0 f	2.4 ab	13.9 a
Mean	5.8	4.8	0.9	3.5	80.7	2.2	12.0
LSD¹	0.1	0.1	0.1	0.5	1.5	0.4	1.2

¹ Means sharing the same letter(s) are not statistically different, at P=0.05 based on the Fisher's LSD test.

Fatty Acid Results

Table 20. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk, VA), Dig 1, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.2 a	2.3 d-h	53.6 m	28.3 a	1.2 b-e	1.4 k
Bailey II	6.0 l	2.3 d-g	80.3 ab	5.2 jk	1.1 f-j	1.7 d-j
Emery	6.0 l	2.4 c-e	79.8 a-c	5.5 jk	1.1 d-g	1.7 c-j
Sullivan	6.1 j-l	2.3 e-j	78.9 a-e	6.2 h-k	1.1 e-i	1.8 c-j
Walton	6.1 kl	2.3 d-h	76.8 c-f	7.4 h-j	1.2 c-g	2.1 a
Wynne	6.8 g-i	2.4 c-f	73.9 f-i	10.5 e-g	1.2 c-f	1.7 f-j
N14001	6.0 l	2.4 c-e	81.0 a	4.7 jk	1.1 f-j	1.7 g-j
N14002oIJ	6.7 g-i	2.3 e-k	76.1 e-h	8.4 g-i	1.1 f-j	1.8 c-j
N14004oIJ	7.1 e-g	2.4 c-e	72.0 ij	12.0 de	1.2 c-f	1.7 h-j
N14007	8.1 b	2.0 m	64.3 l	19.2 b	1.0 k-n	1.7 g-j
N14009	5.9 l	2.1 h-m	79.6 a-c	5.8 i-k	1.1 i-n	1.9 c-h
N15066	6.9 f-h	2.1 k-m	75.8 e-h	8.6 f-h	1.0 l-n	1.9 b-d
N14023oI	6.2 j-l	2.0 lm	80.2 ab	5.2 jk	1.0 n	1.9 b-g
N14027oIJ	7.8 bc	2.2 g-m	67.2 kl	16.6 bc	1.1 h-n	1.6 ij
N15017oI	6.4 i-k	2.2 e-l	77.6 b-e	7.3 h-j	1.1 f-l	1.9 b-g
N15039oI	6.7 g-i	2.3 d-i	74.2 f-i	10.4 e-g	1.1 e-h	1.7 e-j
N15041oI	7.1 e-g	2.2 g-m	73.3 g-i	11.5 e	1.1 j-n	1.6 j
N15044oIF	7.5 cd	2.1 j-m	69.1 jk	15.1 c	1.1 i-n	1.7 g-j
N15053	6.5 h-j	2.1 i-m	76.2 d-g	8.4 g-i	1.1 g-m	1.9 c-g
N15060	5.9 l	2.2 e-l	78.7 a-e	6.0 h-k	1.1 f-l	2.0 ab
N16005	7.2 d-f	2.7 b	73.0 hi	11.2 ef	1.2 b	1.3 k
N16012	6.2 j-l	2.2 e-k	78.4 a-e	6.9 h-j	1.1 f-k	1.7 d-j
N16021	7.4 c-e	2.1 g-m	69.2 jk	14.6 cd	1.1 f-k	1.8 c-j
N17036	6.1 j-l	2.5 b-d	79.4 a-d	5.4 jk	1.2 bc	1.8 c-i
N17037	6.2 j-l	2.5 bc	78.9 a-e	5.8 i-k	1.2 b-d	1.7 d-j
N17040	6.1 kl	2.0 lm	79.4 a-d	6.1 h-k	1.0 n	1.9 b-f
N17041	6.1 kl	2.1 k-m	79.6 a-c	5.9 h-k	1.0 mn	1.9 b-e
N17044	5.8 l	3.1 a	80.0 a-c	4.0 k	1.4 a	1.6 ij
N17045	6.4 i-k	2.1 k-m	78.1 a-e	7.0 h-j	1.0 mn	1.9 a-c
N17047	6.0 l	2.2 f-m	80.6 ab	4.9 jk	1.1 j-n	1.9 c-h
Mean	6.6	2.3	75.5	9.1	1.1	1.8
LSD²	0.4	0.2	3.2	2.8	0.1	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 20. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk, VA) Dig 1, 2020¹, (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.7 a	1.2 c-h	96.3 a	1.9 l	16.6 a	1.7 a	5.1 b-d
Bailey II	2.2 e-g	1.2 c-g	79.4 ij	15.9 b-d	12.8 kl	0.4 i-k	4.5 i-l
Emery	2.4 c-e	1.1 f-h	79.5 ij	14.6 b-f	13.0 i-l	0.4 i-k	4.6 e-j
Sullivan	2.3 c-g	1.2 c-g	80.1 g-j	12.7 d-h	13.0 h-l	0.5 g-j	4.6 e-j
Walton	2.7 a	1.4 a	80.5 g-j	10.6 g-i	13.7 e-g	0.5 g-j	5.3 ab
Wynne	2.5 b-d	1.1 d-h	83.1 ef	7.0 i-k	13.9 d-f	0.8 ef	4.8 e-h
N14001	2.1 fg	1.0 h	79.2 j	17.3 ab	12.6 l	0.4 jk	4.2 l
N14002oIJ	2.4 b-d	1.2 c-g	81.4 f-i	9.1 h-j	13.7 e-g	0.6 f-h	4.8 e-h
N14004oIJ	2.5 b-d	1.3 a-e	84.0 e	6.1 jk	14.3 b-d	0.8 e	4.9 c-e
N14007	2.5 b-d	1.2 b-f	89.9 b	3.4 kl	14.8 b	1.3 b	4.7 e-i
N14009	2.4 b-d	1.2 b-f	79.9 g-j	14.4 b-g	12.7 kl	0.5 h-j	4.7 e-h
N15066	2.5 b-d	1.2 b-f	81.6 f-h	8.9 h-j	13.7 e-g	0.6 fg	4.7 e-i
N14023oI	2.2 e-g	1.2 a-f	79.6 ij	15.3 b-e	12.7 l	0.4 i-k	4.4 i-l
N14027oIJ	2.3 c-f	1.2 c-g	87.9 bc	4.0 kl	14.5 bc	1.1 bc	4.6 e-k
N15017oI	2.4 c-e	1.2 c-h	80.8 g-j	11.1 f-h	13.3 g-k	0.5 g-i	4.6 e-j
N15039oI	2.3 c-f	1.2 c-g	83.2 ef	7.1 i-k	13.7 e-g	0.8 ef	4.7 e-j
N15041oI	2.1 fg	1.2 c-h	84.3 de	6.8 i-k	13.5 e-i	0.8 de	4.3 kl
N15044oIF	2.1 g	1.3 a-f	87.0 c	4.6 kl	14.1 c-e	1.1 c	4.4 j-l
N15053	2.4 cd	1.3 a-c	81.7 fg	9.1 h-j	13.5 f-j	0.6 fg	4.8 d-g
N15060	2.7 ab	1.4 ab	79.6 h-j	13.3 c-g	13.3 g-k	0.4 h-j	5.1 bc
N16005	2.3 c-g	1.1 gh	83.3 ef	6.6 jk	14.5 b-d	0.8 ef	4.6 f-k
N16012	2.3 c-e	1.1 e-h	80.7 g-j	11.7 e-h	13.0 h-l	0.5 g-j	4.6 f-k
N16021	2.5 bc	1.2 b-f	86.2 cd	4.8 kl	14.4 b-d	1.0 cd	4.8 c-f
N17036	2.4 b-d	1.2 c-g	79.0 jk	14.9 b-f	13.5 f-j	0.4 i-k	4.8 c-f
N17037	2.5 bc	1.1 d-h	79.2 j	13.7 b-g	13.6 e-h	0.4 i-k	4.8 d-f
N17040	2.3 c-g	1.2 c-f	80.3 g-j	13.0 c-g	12.6 l	0.5 g-j	4.5 g-l
N17041	2.3 c-g	1.2 c-f	80.1 g-j	13.5 b-g	12.6 l	0.5 g-j	4.5 g-l
N17044	2.8 a	1.3 a-d	77.0 k	20.3 a	14.4 b-d	0.3 k	5.5 a
N17045	2.2 e-g	1.2 c-g	80.9 g-j	11.2 f-h	12.9 j-l	0.5 g-j	4.4 j-l
N17047	2.2 d-g	1.2 c-g	79.3 j	16.5 a-c	12.6 l	0.4 i-k	4.5 h-l
Mean	2.4	1.2	82.2	10.6	13.6	0.7	4.7
LSD²	0.2	0.1	2.0	3.8	0.6	0.2	0.3

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 21. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk, VA), Dig 2, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	8.4 a	1.9 cd	58.6 c	25.1 a	1.0 cd	1.4 d
Bailey II	5.9 c-e	2.2 a-d	79.4 a	6.3 c	1.1 a-c	1.8 a-c
Emery	5.7 c-e	2.4 a-c	81.4 a	4.4 c	1.1 a-c	1.7 a-d
Sullivan	5.8 c-e	2.0 cd	80.4 a	5.8 c	1.0 cd	1.7 a-c
Walton	5.1 e	1.9 d	81.8 a	5.6 c	0.8 d	1.7 a-c
Wynne	5.7 c-e	2.6 a	80.2 a	5.2 c	1.2 a	1.7 a-d
N14001	6.5 b-d	2.3 a-d	75.8 ab	9.5 bc	1.1 a-c	1.6 b-d
N14002oIJ	5.9 c-e	2.1 b-d	80.7 a	5.2 c	1.0 b-d	1.8 a-c
N14004oIJ	5.7 c-e	2.2 a-d	81.2 a	5.2 c	1.0 b-d	1.6 b-d
N14007	6.1 c-e	2.1 b-d	78.5 a	7.4 c	1.0 b-d	1.7 a-d
N14009	5.4 de	2.1 b-d	81.1 a	5.0 c	1.0 b-d	2.0 a
N15066	6.2 c-e	2.0 cd	78.4 a	7.2 c	1.0 cd	1.8 ab
N14023oI	5.7 c-e	2.2 a-d	81.4 a	4.6 c	1.1 a-c	1.7 a-c
N14027oIJ	6.4 b-e	2.0 cd	75.7 ab	9.9 bc	1.0 b-d	1.7 a-d
N15017oI	6.2 c-e	2.3 a-d	77.7 a	7.6 c	1.1 a-c	1.7 a-d
N15039oI	6.2 c-e	2.2 a-d	77.6 a	7.9 c	1.1 a-c	1.7 a-d
N15041oI	6.5 b-d	2.0 cd	76.6 a	9.2 c	1.0 cd	1.6 a-d
N15044oIF	6.6 b-d	2.0 b-d	74.3 ab	11.0 bc	1.0 b-d	1.6 b-d
N15053	5.9 c-e	2.0 b-d	78.4 a	7.3 c	1.0 b-d	1.9 ab
N15060	6.0 c-e	2.0 b-d	77.2 a	8.2 c	1.0 b-d	1.9 ab
N16005	7.0 bc	2.2 a-d	71.8 ab	13.2 bc	1.1 a-c	1.5 b-d
N16012	5.6 de	2.1 b-d	81.9 a	4.4 c	1.0 b-d	1.8 a-c
N16021	6.5 b-d	2.1 b-d	74.6 ab	10.6 bc	1.1 a-c	1.7 a-d
N17036	7.7 ab	2.2 a-d	65.3 bc	18.7 ab	1.1 a-c	1.5 cd
N17037	6.1 c-e	2.3 a-d	78.1 a	7.1 c	1.1 a-c	1.8 a-c
N17040	6.0 c-e	2.2 a-d	78.3 a	7.5 c	1.1 a-c	1.7 a-d
N17041	6.0 c-e	2.0 cd	79.4 a	6.7 c	0.9 cd	1.7 a-c
N17044	5.6 de	2.5 ab	81.3 a	4.4 c	1.2 ab	1.7 a-d
N17045	5.9 c-e	2.2 a-d	79.9 a	6.0 c	1.0 b-d	1.7 a-d
N17047	5.7 de	2.2 a-d	81.1 a	5.0 c	1.1 a-c	1.8 a-c
Mean	6.1	2.2	77.6	8.0	1.0	1.7
LSD²	1.3	0.5	10.6	9.3	0.2	0.4

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 21. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Tidewater AREC (Suffolk, VA), Dig 2, 2020¹ (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.3 a	1.3 a	95.0 a	2.6 f	14.9 a	1.7 a	4.6 a-c
Bailey II	2.1 a-e	1.2 ab	80.7 c	12.6 a-e	12.5 cd	0.5 cd	4.3 a-d
Emery	2.1 a-e	1.2 ab	79.0 c	18.5 ab	12.5 cd	0.4 d	4.4 a-c
Sullivan	2.1 a-e	1.2 ab	80.6 c	14.5 a-e	12.1 cd	0.5 cd	4.3 a-d
Walton	2.0 b-e	1.1 b	81.4 c	15.4 a-d	10.9 d	0.5 cd	3.9 d
Wynne	2.2 a-e	1.3 a	79.3 c	15.5 a-d	13.0 bc	0.4 cd	4.7 a
N14001	2.0 a-e	1.2 ab	82.9 bc	8.2 c-f	13.1 bc	0.7 b-d	4.3 a-d
N14002oIJ	2.0 a-e	1.3 a	79.9 c	15.9 a-d	12.3 cd	0.4 cd	4.3 a-d
N14004oIJ	1.9 e	1.2 ab	80.1 c	15.7 a-d	12.0 cd	0.4 cd	4.1 cd
N14007	2.0 b-e	1.2 ab	81.6 c	10.8 a-f	12.5 cd	0.6 cd	4.2 b-d
N14009	2.2 a-e	1.3 a	79.9 c	16.3 a-d	12.0 cd	0.4 cd	4.5 a-c
N15066	2.0 a-e	1.3 a	81.5 c	11.5 a-f	12.5 cd	0.6 cd	4.3 a-d
N14023oI	2.0 a-e	1.2 ab	79.4 c	17.6 a-c	12.2 cd	0.4 d	4.3 a-d
N14027oIJ	2.0 b-e	1.3 a	83.6 bc	10.7 a-f	12.7 c	0.8 b-d	4.2 a-d
N15017oI	2.1 a-e	1.3 a	81.4 c	10.9 a-f	13.0 bc	0.6 cd	4.5 a-c
N15039oI	2.2 a-e	1.2 ab	81.7 c	12.1 a-f	12.9 c	0.6 cd	4.5 a-c
N15041oI	1.9 de	1.3 a	83.0 bc	9.0 b-f	12.6 c	0.7 b-d	4.2 b-d
N15044oIF	2.1 a-e	1.2 ab	84.3 bc	8.0 c-f	13.0 bc	0.8 b-d	4.4 a-d
N15053	2.2 a-e	1.3 a	81.6 c	10.9 a-f	12.4 cd	0.6 cd	4.5 a-c
N15060	2.3 ab	1.3 a	82.1 c	11.3 a-f	12.7 c	0.6 b-d	4.6 ab
N16005	2.1 a-e	1.2 ab	85.8 bc	5.5 ef	13.5 a-c	1.0 bc	4.3 a-d
N16012	2.0 b-e	1.2 ab	79.5 c	18.8 a	11.9 cd	0.4 d	4.2 b-d
N16021	2.2 a-e	1.3 a	83.8 bc	9.6 a-f	13.1 bc	0.8 b-d	4.5 a-c
N17036	2.3 a-c	1.3 a	89.7 ab	6.6 d-f	14.6 ab	1.2 ab	4.6 ab
N17037	2.2 a-d	1.3 a	80.9 c	12.7 a-e	13.0 bc	0.5 cd	4.6 a-c
N17040	2.1 a-e	1.2 ab	81.5 c	12.3 a-e	12.6 c	0.6 cd	4.3 a-d
N17041	2.0 c-e	1.3 a	81.3 c	13.5 a-e	12.1 cd	0.5 cd	4.1 b-d
N17044	2.2 a-e	1.2 ab	78.9 c	18.9 a	12.7 c	0.4 d	4.6 a-c
N17045	2.1 a-e	1.3 a	80.3 c	14.7 a-e	12.5 cd	0.5 cd	4.4 a-d
N17047	2.0 b-e	1.2 ab	79.8 c	16.5 a-c	12.1 cd	0.4 cd	4.3 a-d
Mean	2.1	1.2	82.0	12.6	12.7	0.6	4.4
LSD²	0.3	0.2	7.1	9.7	1.7	0.6	0.5

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 22. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Averages of all Digs from Tidewater AREC (Suffolk, VA), 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	8.8 a	2.1 c-i	56.1 g	26.7 a	1.1 b-g	1.4 h
Bailey II	5.9 e-g	2.3 b-h	79.9 a	5.8 gh	1.1 b-g	1.7 b-g
Emery	5.8 fg	2.4 b-e	80.6 a	5.0 h	1.1 b-e	1.7 c-g
Sullivan	6.0 e-g	2.2 c-i	79.6 a	6.0 gh	1.1 c-g	1.8 a-g
Walton	5.6 g	2.1 f-i	79.3 a	6.5 gh	1.0 fg	1.9 ab
Wynne	6.2 c-g	2.5 b	77.0 a-f	7.8 d-h	1.2 ab	1.7 d-g
N14001	6.2 d-g	2.3 b-g	78.4 ab	7.1 e-h	1.1 b-f	1.6 g
N14002oIJ	6.3 b-g	2.2 c-i	78.4 a-c	6.8 f-h	1.1 c-g	1.8 a-g
N14004oIJ	6.4 b-g	2.3 b-i	76.6 a-f	8.6 b-h	1.1 b-g	1.6 fg
N14007	7.1 b	2.1 hi	71.4 f	13.3 b	1.0 d-g	1.7 e-g
N14009	5.6 g	2.1 e-i	80.3 a	5.4 gh	1.0 c-g	1.9 ab
N15066	6.5 b-f	2.0 hi	77.1 a-f	7.9 c-h	1.0 fg	1.9 a-c
N14023oI	5.9 e-g	2.1 e-i	80.8 a	4.9 h	1.0 c-g	1.8 a-g
N14027oIJ	7.1 b	2.1 g-i	71.5 f	13.3 b	1.0 c-g	1.7 d-g
N15017oI	6.3 b-g	2.2 b-i	77.6 a-e	7.5 d-h	1.1 b-g	1.8 a-g
N15039oI	6.4 b-g	2.3 b-i	75.9 a-f	9.2 b-h	1.1 b-f	1.7 c-g
N15041oI	6.8 b-e	2.1 g-i	74.9 a-f	10.4 b-g	1.0 e-g	1.6 fg
N15044oIF	7.1 bc	2.1 g-i	71.7 ef	13.1 bc	1.0 c-g	1.6 e-g
N15053	6.2 d-g	2.1 g-i	77.3 a-f	7.9 c-h	1.0 c-g	1.9 a-d
N15060	6.0 e-g	2.1 d-i	78.0 a-d	7.1 e-h	1.1 c-g	2.0 a
N16005	7.1 b	2.4 bc	72.4 b-f	12.2 b-e	1.1 b-d	1.4 h
N16012	5.9 fg	2.2 c-i	80.2 a	5.6 gh	1.1 c-g	1.7 b-g
N16021	7.0 b-d	2.1 e-i	71.9 d-f	12.6 b-d	1.1 b-g	1.7 c-g
N17036	6.9 b-d	2.4 b-f	72.3 c-f	12.0 b-f	1.2 bc	1.7 e-g
N17037	6.2 d-g	2.4 b-d	78.5 ab	6.5 gh	1.2 bc	1.8 a-g
N17040	6.1 e-g	2.1 e-i	78.8 a	6.8 gh	1.0 c-g	1.8 a-g
N17041	6.0 e-g	2.0 i	79.5 a	6.3 gh	1.0 g	1.8 a-g
N17044	5.7 fg	2.8 a	80.6 a	4.2 h	1.3 a	1.6 e-g
N17045	6.2 d-g	2.1 e-i	79.0 a	6.5 gh	1.0 c-g	1.8 a-f
N17047	5.8 fg	2.2 c-i	80.8 a	4.9 h	1.1 c-g	1.8 a-e
Mean	6.4	2.2	76.6	8.6	1.1	1.7
LSD²	0.8	0.3	6.1	5.2	0.1	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 22. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Average of all Digs from Tidewater AREC (Suffolk, VA), 2020¹ (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.5 a	1.2 b-e	95.6 a	2.2 k	15.8 a	1.7 a	4.8 a-c
Bailey II	2.1 cd	1.2 b-e	80.0 f-h	14.2 a-e	12.6 c-g	0.5 gh	4.4 d-f
Emery	2.2 a-d	1.2 c-e	79.2 gh	16.6 ab	12.7 c-g	0.4 gh	4.5 b-f
Sullivan	2.2 b-d	1.2 b-e	80.3 f-h	13.6 b-e	12.6 c-g	0.5 f-h	4.4 c-f
Walton	2.3 a-c	1.2 a-e	81.0 d-h	13.0 b-g	12.3 g	0.5 f-h	4.6 b-f
Wynne	2.3 a-c	1.2 b-e	81.2 c-h	11.3 b-j	13.4 b-g	0.6 d-h	4.7 a-d
N14001	2.1 cd	1.1 e	81.0 d-h	12.7 b-h	12.8 b-g	0.5 f-h	4.3 ef
N14002oIJ	2.2 a-d	1.2 a-e	80.7 e-h	12.5 b-i	13.0 b-g	0.5 f-h	4.5 b-f
N14004oIJ	2.2 b-d	1.2 a-e	82.1 b-g	10.9 c-j	13.2 b-g	0.6 c-h	4.5 b-f
N14007	2.2 a-d	1.2 b-e	85.7 b	7.1 i-k	13.7 b-d	0.9 bc	4.5 b-f
N14009	2.3 a-d	1.3 a-c	79.9 f-h	15.3 a-d	12.3 fg	0.4 gh	4.6 a-f
N15066	2.2 a-d	1.3 a-c	81.5 c-h	10.2 d-j	13.1 b-g	0.6 d-h	4.5 b-f
N14023oI	2.1 cd	1.2 a-d	79.5 gh	16.5 a-c	12.5 d-g	0.4 gh	4.4 d-f
N14027oIJ	2.1 cd	1.2 a-e	85.8 b	7.4 g-k	13.6 b-e	1.0 bc	4.4 c-f
N15017oI	2.2 a-d	1.2 a-e	81.1 d-h	11.0 b-j	13.1 b-g	0.6 e-h	4.6 b-f
N15039oI	2.3 a-d	1.2 b-e	82.5 b-g	9.6 e-j	13.3 b-g	0.7 b-g	4.6 b-f
N15041oI	2.0 d	1.2 b-e	83.7 b-f	7.9 f-j	13.1 b-g	0.8 b-f	4.2 f
N15044oIF	2.1 cd	1.2 a-e	85.7 b	6.3 jk	13.5 b-f	1.0 b	4.4 d-f
N15053	2.3 a-d	1.3 ab	81.6 c-h	10.0 d-j	12.9 b-g	0.6 d-h	4.7 a-f
N15060	2.5 ab	1.3 a	80.9 d-h	12.3 b-i	13.0 b-g	0.5 f-h	4.9 ab
N16005	2.2 cd	1.1 de	84.5 b-d	6.0 jk	14.0 b	0.9 b-e	4.4 c-f
N16012	2.2 cd	1.2 c-e	80.1 f-h	15.2 a-e	12.5 d-g	0.4 gh	4.4 d-f
N16021	2.3 a-c	1.3 a-d	85.0 bc	7.2 h-k	13.8 bc	0.9 b-d	4.7 a-e
N17036	2.4 a-c	1.2 a-e	84.3 b-e	10.8 d-j	14.0 b	0.8 b-f	4.7 a-d
N17037	2.4 a-c	1.2 b-e	80.1 f-h	13.2 b-f	13.3 b-g	0.5 f-h	4.7 a-d
N17040	2.2 cd	1.2 b-e	80.9 d-h	12.7 b-i	12.6 c-g	0.5 f-h	4.4 c-f
N17041	2.1 cd	1.2 a-e	80.7 d-h	13.5 b-f	12.4 e-g	0.5 f-h	4.3 d-f
N17044	2.5 a	1.2 a-e	77.9 h	19.6 a	13.5 b-f	0.3 h	5.0 a
N17045	2.1 cd	1.2 a-e	80.6 e-h	12.9 b-g	12.7 c-g	0.5 f-h	4.4 d-f
N17047	2.1 cd	1.2 b-e	79.5 gh	16.5 a-c	12.4 e-g	0.4 gh	4.4 d-f
Mean	2.2	1.2	82.1	11.6	13.1	0.6	4.5
LSD²	0.3	0.1	3.9	5.6	1.2	0.3	0.4

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 23. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC Dig 1, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.4 a	2.4 d-j	53.8 e	28.6 a	1.1 c-f	1.2 m
Bailey II	6.0 j-n	2.4 c-i	80.3 a	5.3 fg	1.1 e-i	1.6 c-g
Emery	5.8 l-n	2.6 bc	81.4 a	4.1 fg	1.2 b-d	1.5 h-k
Sullivan	6.1 i-n	2.2 h-m	80.0 a	5.7 fg	1.1 g-l	1.6 d-h
Walton	5.9 k-n	2.4 c-i	79.8 a	5.0 fg	1.1 c-f	1.9 b
Wynne	6.5 f-k	2.4 c-h	78.1 ab	6.8 fg	1.1 c-f	1.6 f-i
N14001	6.0 j-n	2.4 c-f	81.3 a	4.5 fg	1.1 d-g	1.5 i-l
N14002oIJ	6.5 f-j	2.4 c-e	79.0 a	6.2 fg	1.1 d-h	1.5 i-l
N14004oIJ	7.6 bc	2.3 e-l	70.0 cd	14.2 bc	1.1 f-k	1.4 kl
N14007	8.0 b	2.2 g-m	66.2 d	17.7 b	1.1 e-k	1.4 l
N14009	5.6 n	2.3 e-l	81.9 a	4.2 fg	1.1 e-i	1.6 c-g
N15066	6.8 f-h	2.2 j-n	77.6 ab	7.1 e-g	1.0 h-m	1.7 c
N14023oI	6.2 h-m	2.0 no	81.2 a	4.6 fg	1.0 n	1.7 c-f
N14027oIJ	7.4 b-d	2.1 k-n	70.9 c	13.7 c	1.1 g-l	1.5 j-l
N15017oI	6.2 i-n	2.3 e-l	80.7 a	4.8 fg	1.1 e-i	1.6 d-h
N15039oI	6.9 e-g	2.3 d-k	73.9 bc	11.0 cd	1.1 d-h	1.4 kl
N15041oI	6.6 f-i	2.1 m-o	78.1 ab	7.3 d-f	1.0 l-n	1.6 c-g
N15044oIF	7.3 c-e	2.1 k-n	71.9 c	12.6 c	1.0 g-l	1.5 g-j
N15053	6.4 g-l	2.2 e-m	78.3 a	6.9 fg	1.1 e-j	1.6 d-h
N15060	6.3 h-m	1.9 o	78.7 a	5.8 fg	1.0 mn	2.1 a
N16005	7.3 c-e	2.7 b	72.8 c	11.6 c	1.2 b	1.2 m
N16012	5.9 k-n	2.4 c-g	81.3 a	4.3 fg	1.1 b-e	1.6 f-i
N16021	7.0 d-f	2.2 f-m	73.7 bc	10.8 c-e	1.1 d-i	1.6 g-j
N17036	6.1 i-n	2.5 cd	80.8 a	4.4 fg	1.2 b-d	1.6 e-h
N17037	6.1 i-n	2.5 cd	80.4 a	4.8 fg	1.2 bc	1.6 g-j
N17040	5.9 k-n	2.1 l-o	81.1 a	4.8 fg	1.0 k-n	1.7 c
N17041	6.0 j-n	2.2 i-n	80.9 a	4.8 fg	1.0 i-m	1.7 c-e
N17044	5.8 mn	3.0 a	81.6 a	3.4 g	1.3 a	1.5 h-k
N17045	6.2 i-n	2.1 k-n	80.5 a	5.2 fg	1.0 j-n	1.7 cd
N17047	6.0 i-n	2.1 l-o	81.3 a	4.4 fg	1.0 l-n	1.7 c
Mean	6.5	2.3	77.3	7.8	1.1	1.6
LSD²	0.6	0.2	4.4	3.8	0.1	0.1

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 23. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC Dig 1, 2020¹, (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.3 bc	1.3 c-f	96.7 a	1.9 m	16.4 a	1.7 a	4.7 cd
Bailey II	2.1 g-k	1.3 c-f	79.5 fg	15.3 c-g	12.7 h-l	0.4 e-g	4.4 f-k
Emery	2.1 f-k	1.3 c-f	78.4 fg	19.7 b	12.9 f-l	0.3 fg	4.5 d-j
Sullivan	2.0 g-k	1.3 c-f	80.0 fg	14.0 d-h	12.7 i-l	0.5 e-g	4.4 g-k
Walton	2.4 b	1.5 b	78.8 fg	16.0 b-f	13.3 d-j	0.4 e-g	5.0 ab
Wynne	2.2 c-f	1.3 c-e	80.3 f	11.4 g-i	13.5 d-h	0.5 ef	4.6 c-f
N14001	2.0 i-k	1.2 fg	78.9 fg	17.9 b-e	12.7 i-l	0.4 e-g	4.3 k
N14002oIJ	2.0 g-k	1.3 c-f	79.8 fg	12.8 f-h	13.4 d-i	0.5 e-g	4.4 f-k
N14004oIJ	2.2 d-h	1.3 c-f	85.9 bc	5.1 k-m	14.4 bc	1.0 bc	4.5 d-j
N14007	2.2 c-g	1.3 c-e	88.7 b	3.8 lm	14.7 b	1.2 b	4.6 c-h
N14009	2.1 f-j	1.2 e-g	79.0 fg	19.9 b	12.3 l	0.3 e-g	4.4 g-k
N15066	2.2 b-e	1.4 bc	80.4 f	11.0 g-j	13.5 d-g	0.5 ef	4.6 c-f
N14023oI	1.9 k	1.3 cd	79.1 fg	18.1 b-d	12.5 j-l	0.4 e-g	4.3 jk
N14027oIJ	2.1 e-i	1.3 c-f	85.8 c	5.4 k-m	14.0 b-d	1.0 bc	4.4 e-k
N15017oI	2.0 h-k	1.4 b-d	78.9 fg	17.0 b-f	12.9 f-l	0.4 e-g	4.5 d-k
N15039oI	2.1 e-j	1.3 c-f	83.7 cd	8.3 i-k	13.7 c-f	0.8 cd	4.5 d-k
N15041oI	2.0 jk	1.3 c-e	81.1 d-f	10.8 h-j	12.9 f-l	0.6 de	4.3 i-k
N15044oIF	2.1 f-j	1.3 c-e	84.9 c	5.7 k-m	13.9 c-e	0.9 c	4.5 d-k
N15053	2.2 c-h	1.3 c-e	80.6 ef	11.4 g-i	13.2 e-k	0.5 ef	4.5 c-i
N15060	2.6 a	1.6 a	79.5 fg	13.6 e-h	13.4 d-i	0.4 e-g	5.1 a
N16005	2.0 i-k	1.1 g	83.6 cd	7.1 i-l	14.4 bc	0.8 cd	4.3 h-k
N16012	2.2 c-h	1.2 d-g	78.6 fg	19.1 bc	12.9 g-l	0.3 e-g	4.5 c-i
N16021	2.3 b-d	1.3 c-e	83.4 c-e	6.8 j-l	13.9 c-e	0.8 cd	4.7 c-e
N17036	2.2 c-h	1.3 c-f	78.4 fg	18.6 bc	13.2 e-j	0.3 e-g	4.6 c-h
N17037	2.2 c-g	1.3 c-f	78.7 fg	16.8 b-f	13.3 d-j	0.4 e-g	4.6 c-g
N17040	2.1 g-k	1.3 c-f	79.4 fg	16.9 b-f	12.4 kl	0.4 e-g	4.3 h-k
N17041	2.1 f-k	1.3 c-e	79.3 fg	16.8 b-f	12.5 j-l	0.4 e-g	4.4 f-k
N17044	2.3 b-d	1.2 e-g	77.2 g	24.4 a	13.5 d-g	0.2 g	4.8 bc
N17045	2.0 i-k	1.3 c-e	79.5 fg	15.9 b-f	12.6 i-l	0.4 e-g	4.3 h-k
N17047	2.1 g-k	1.4 cd	78.9 fg	18.4 bc	12.6 j-l	0.4 e-g	4.4 f-k
Mean	2.1	1.3	81.2	13.3	13.3	0.6	4.5
LSD²	0.1	0.1	2.8	4.4	0.8	0.2	0.3

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 24. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC Dig 2, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.4 a	2.3 b-g	54.1 l	28.3 a	1.1 b-d	1.2 kl
Bailey II	5.9 g-i	2.3 b-f	80.5 a-d	5.5 h-k	1.0 b-f	1.6 b-h
Emery	5.9 f-i	2.4 b-e	80.9 a-d	5.0 i-k	1.1 b-e	1.6 c-i
Sullivan	6.1 e-h	2.2 d-i	79.9 a-e	6.0 g-j	1.0 b-f	1.6 b-g
Walton	5.9 g-i	2.2 c-h	79.6 b-f	6.0 g-j	1.0 b-f	1.7 a-c
Wynne	6.3 e-g	2.4 b-e	79.6 c-f	6.2 f-j	1.1 b-e	1.5 e-j
N14001	5.8 g-i	2.3 b-g	82.7 a	3.7 jk	1.0 b-f	1.5 f-j
N14002oIJ	6.2 e-g	2.4 b-d	80.7 a-d	4.9 i-k	1.1 b-e	1.5 d-i
N14004oIJ	6.0 f-i	1.9 i	74.3 h-j	13.3 cd	0.8 g	1.1 l
N14007	8.1 b	2.2 d-h	63.0 k	20.9 b	1.1 b-e	1.3 i-l
N14009	5.7 hi	2.2 d-i	82.2 a-c	4.3 i-k	1.0 c-f	1.6 b-h
N15066	6.9 cd	1.9 hi	77.1 e-h	8.4 e-h	0.9 fg	1.6 b-h
N14023oI	5.9 f-i	2.1 e-i	82.3 a-c	4.1 i-k	1.0 d-g	1.6 b-h
N14027oIJ	7.3 c	2.1 d-i	71.5 j	13.7 c	1.0 c-f	1.4 h-k
N15017oI	6.2 e-g	2.0 g-i	79.6 c-f	6.4 f-j	1.0 c-g	1.7 a-d
N15039oI	6.6 de	2.4 b-e	75.0 hi	10.2 de	1.1 b-d	1.4 g-k
N15041oI	6.6 de	2.0 f-i	78.3 d-g	7.1 f-i	1.0 e-g	1.6 b-h
N15044oIF	7.3 c	2.1 e-i	72.6 ij	12.4 cd	1.0 d-g	1.6 b-h
N15053	6.5 de	2.3 b-g	76.8 f-h	8.6 e-h	1.1 b-e	1.5 c-i
N15060	5.7 g-i	2.0 g-i	81.1 a-d	4.8 i-k	1.0 d-g	1.9 a
N16005	6.9 cd	2.6 b	76.0 gh	9.2 ef	1.2 b	1.2 j-l
N16012	6.0 f-i	2.5 bc	81.0 a-d	4.4 i-k	1.2 bc	1.5 c-i
N16021	6.5 d-f	2.2 c-h	76.5 gh	8.9 e-g	1.1 b-e	1.6 b-h
N17036	5.9 g-i	2.2 d-h	80.7 a-d	5.0 i-k	1.1 b-f	1.7 a-d
N17037	5.9 g-i	2.6 b	81.2 a-d	4.4 i-k	1.2 b	1.6 b-h
N17040	5.8 g-i	2.1 d-i	81.3 a-d	4.8 i-k	1.0 b-f	1.7 a-c
N17041	5.7 g-i	2.1 d-i	81.8 a-c	4.4 i-k	1.0 b-f	1.7 b-f
N17044	5.5 i	3.0 a	82.6 ab	2.8 k	1.3 a	1.5 e-j
N17045	6.2 e-g	2.0 f-i	80.5 a-d	5.7 h-k	1.0 e-g	1.7 a-e
N17047	6.0 f-i	2.0 g-i	81.2 a-d	4.7 i-k	1.0 d-g	1.8 ab
Mean	6.4	2.2	77.8	7.8	1.0	1.5
LSD²	0.5	0.3	3.0	3.1	0.2	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 24. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Martin County, NC Dig 2, 2020¹, (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.3 a	1.1 b-d	96.6 a	1.9 p	16.3 a	1.7 a	4.6 ab
Bailey II	2.0 c-h	1.2 a-d	79.9 g-k	14.9 e-h	12.5 c-f	0.4 g-l	4.2 a-f
Emery	1.9 d-i	1.2 a-d	79.5 g-k	16.2 d-g	12.6 c-f	0.4 g-l	4.2 a-f
Sullivan	2.0 d-h	1.1 b-d	80.4 f-j	13.8 f-i	12.4 d-g	0.5 f-l	4.1 a-f
Walton	2.1 a-d	1.4 ab	80.3 f-j	13.3 g-i	12.6 c-f	0.5 f-l	4.5 a-c
Wynne	2.0 d-h	1.1 b-d	80.3 f-j	12.9 hi	12.8 c-f	0.5 f-l	4.1 a-f
N14001	1.9 d-i	1.1 b-d	78.7 jk	22.1 b	12.2 fg	0.3 kl	4.0 b-f
N14002oIJ	2.0 c-h	1.2 a-d	79.0 i-k	16.7 d-f	12.9 c-f	0.4 g-l	4.3 a-f
N14004oIJ	1.7 i	0.9 d	87.8 c	6.1 l-o	11.3 g	1.3 bc	3.5 g
N14007	2.2 ab	1.2 a-d	91.4 b	3.1 op	14.8 b	1.4 b	4.5 a-d
N14009	2.0 d-h	1.2 a-d	79.3 h-k	19.2 b-d	12.0 fg	0.4 h-l	4.1 a-f
N15066	2.0 b-g	1.1 b-d	82.2 e-h	9.3 j-l	12.8 c-f	0.7 e-i	4.0 c-f
N14023oI	1.8 hi	1.2 a-d	79.2 h-k	20.1 bc	12.0 fg	0.3 i-l	4.0 e-g
N14027oIJ	1.9 d-i	1.1 b-d	86.2 cd	5.2 no	13.5 cd	1.0 cd	4.0 c-f
N15017oI	2.0 c-h	1.1 b-d	80.9 f-j	12.5 h-j	12.3 d-g	0.5 f-l	4.1 a-f
N15039oI	2.1 b-f	1.2 a-d	83.3 d-f	7.3 l-n	13.4 c-e	0.8 d-f	4.4 a-f
N15041oI	1.9 e-i	1.5 a	80.9 f-j	11.4 i-k	13.0 c-f	0.5 f-k	4.3 a-f
N15044oIF	1.8 g-i	1.1 a-d	85.2 c-e	6.0 m-o	13.4 c-e	0.9 de	4.0 d-g
N15053	2.0 b-g	1.2 a-d	82.1 f-i	9.0 k-m	13.1 c-f	0.7 e-j	4.3 a-f
N15060	2.1 a-d	1.4 ab	79.6 g-k	16.9 c-f	12.2 e-g	0.4 g-l	4.5 a-e
N16005	2.0 b-g	1.0 cd	82.2 e-h	8.3 k-n	13.6 bc	0.7 e-h	4.1 a-f
N16012	2.1 a-d	1.3 a-c	78.5 jk	18.3 cd	13.1 c-f	0.3 j-l	4.5 a-c
N16021	2.1 b-f	1.2 a-c	82.4 e-g	8.8 k-m	13.1 c-f	0.7 e-g	4.4 a-f
N17036	2.2 a-c	1.2 a-c	79.5 g-k	16.2 d-g	12.6 c-f	0.4 g-l	4.5 a-d
N17037	2.1 b-e	1.2 a-d	78.6 jk	18.6 cd	12.9 c-f	0.3 j-l	4.5 a-e
N17040	2.0 d-h	1.2 a-d	79.7 g-k	16.8 d-f	12.1 fg	0.4 g-l	4.2 a-f
N17041	1.9 d-i	1.3 a-c	79.3 h-k	18.6 cd	12.1 fg	0.4 h-l	4.2 a-f
N17044	2.1 a-d	1.1 b-d	77.1 k	29.2 a	13.1 c-f	0.2 i	4.6 a
N17045	1.9 f-i	1.1 b-d	80.4 f-j	14.1 f-i	12.1 fg	0.5 f-l	3.9 fg
N17047	2.1 b-f	1.3 ab	79.3 h-k	17.7 c-e	12.4 d-g	0.4 g-l	4.4 a-f
Mean	2.0	1.2	81.7	13.5	12.8	0.6	4.2
LSD²	0.2	0.3	3.1	3.3	1.2	0.3	0.5

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 25. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Average of Digs from Martin County, NC, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.4 a	2.3 d-h	54.0 k	28.5 a	1.1 b-f	1.2 o
Bailey II	5.9 i-k	2.4 d-g	80.4 a-c	5.4 h-j	1.1 d-j	1.6 d-i
Emery	5.9 jk	2.5 cd	81.2 ab	4.6 i-k	1.1 b-e	1.5 g-l
Sullivan	6.1 h-j	2.2 g-l	79.9 a-d	5.9 g-j	1.1 e-k	1.6 d-h
Walton	5.9 jk	2.3 d-i	79.7 a-e	5.5 h-j	1.1 c-h	1.8 b
Wynne	6.4 f-i	2.4 c-f	78.8 b-e	6.5 g-i	1.1 b-f	1.5 h-l
N14001	5.9 jk	2.4 c-g	82.0 a	4.1 jk	1.1 d-i	1.5 j-m
N14002oIJ	6.4 f-i	2.4 c-e	79.8 a-d	5.5 h-j	1.1 b-g	1.5 i-l
N14004oIJ	6.8 d-f	2.1 k-m	72.2 hi	13.7 c	1.0 l	1.3 no
N14007	8.0 b	2.2 f-l	64.6 j	19.3 b	1.1 d-i	1.4 mn
N14009	5.6 k	2.2 f-k	82.0 a	4.2 jk	1.0 e-l	1.6 e-i
N15066	6.8 de	2.0 lm	77.4 ef	7.8 fg	1.0 kl	1.7 c-f
N14023oI	6.1 h-j	2.0 lm	81.8 a	4.3 jk	1.0 kl	1.7 c-f
N14027oIJ	7.4 c	2.1 i-m	71.2 i	13.7 c	1.0 g-l	1.4 k-m
N15017oI	6.2 g-j	2.1 i-m	80.1 a-c	5.6 h-j	1.0 f-l	1.6 c-g
N15039oI	6.7 d-f	2.4 c-g	74.4 gh	10.6 de	1.1 b-f	1.4 lm
N15041oI	6.6 e-g	2.1 k-m	78.2 c-e	7.2 gh	1.0 kl	1.6 e-i
N15044oIF	7.3 c	2.1 i-m	72.3 hi	12.5 cd	1.0 h-l	1.6 f-k
N15053	6.5 e-h	2.3 e-j	77.6 de	7.7 fg	1.1 c-h	1.6 e-j
N15060	6.0 i-k	2.0 m	79.9 a-d	5.3 h-j	1.0 kl	2.0 a
N16005	7.1 cd	2.7 b	74.4 gh	10.4 e	1.2 b	1.2 o
N16012	6.0 i-k	2.5 cd	81.1 ab	4.4 jk	1.1 b-d	1.6 f-k
N16021	6.7 d-f	2.2 f-l	75.1 fg	9.9 ef	1.1 c-h	1.6 e-j
N17036	6.0 i-k	2.4 d-g	80.7 ab	4.7 i-k	1.1 b-g	1.7 c-f
N17037	6.0 i-k	2.6 bc	80.8 ab	4.6 i-k	1.2 bc	1.6 f-k
N17040	5.9 jk	2.1 i-m	81.2 ab	4.8 i-k	1.0 h-l	1.7 b-d
N17041	5.8 jk	2.2 h-m	81.4 a	4.6 i-k	1.0 g-l	1.7 c-f
N17044	5.7 k	3.0 a	82.1 a	3.1 k	1.3 a	1.5 i-l
N17045	6.2 h-j	2.1 j-m	80.5 a-c	5.4 h-j	1.0 j-l	1.7 c-e
N17047	6.0 i-k	2.1 k-m	81.2 ab	4.5 i-k	1.0 i-l	1.7 bc
Mean	6.4	2.3	77.5	7.8	1.1	1.6
LSD²	0.4	0.2	2.4	2.1	0.1	0.1

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 25. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Average of Digs from Martin County, NC, 2020¹ (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.3 ab	1.2 d-g	96.7 a	1.9 n	16.4 a	1.7 a	4.6 a-d
Bailey II	2.0 e-j	1.2 b-g	79.7 g-i	15.1 e-g	12.6 g-l	0.4 h-j	4.3 d-h
Emery	2.0 f-j	1.2 b-g	78.9 ij	18.0 b-d	12.7 f-l	0.4 jk	4.4 c-g
Sullivan	2.0 f-j	1.2 c-g	80.2 g-i	13.9 g-i	12.6 g-l	0.5 h-j	4.3 d-h
Walton	2.2 a-c	1.4 ab	79.5 g-i	14.6 f-h	12.9 d-l	0.4 h-j	4.7 ab
Wynne	2.1 c-i	1.2 d-g	80.3 g-i	12.1 h-j	13.1 d-i	0.5 h-j	4.4 b-g
N14001	2.0 h-j	1.1 e-g	78.8 ij	20.0 b	12.4 h-l	0.3 jk	4.2 f-h
N14002oIJ	2.0 f-j	1.2 c-g	79.4 hi	14.7 f-h	13.1 d-i	0.4 h-j	4.3 c-h
N14004oIJ	1.9 ij	1.1 fg	86.9 c	5.6 lm	12.8 e-l	1.1 c	4.0 h
N14007	2.2 a-e	1.3 b-f	90.0 b	3.4 mn	14.8 b	1.3 b	4.5 a-f
N14009	2.0 d-j	1.2 d-g	79.2 i	19.6 bc	12.1 l	0.3 jk	4.3 d-h
N15066	2.1 c-h	1.2 b-g	81.3 fg	10.2 jk	13.2 d-h	0.6 gh	4.3 c-h
N14023oI	1.9 j	1.3 b-f	79.1 i	19.1 bc	12.2 kl	0.4 jk	4.1 gh
N14027oIJ	2.0 d-j	1.2 d-g	86.0 c	5.3 lm	13.7 cd	1.0 cd	4.2 e-h
N15017oI	2.0 f-j	1.2 b-g	79.9 g-i	14.7 f-h	12.6 g-l	0.4 h-j	4.3 d-h
N15039oI	2.1 c-i	1.2 c-g	83.5 de	7.8 kl	13.5 c-f	0.8 ef	4.4 b-g
N15041oI	1.9 ij	1.4 a-c	81.0 gh	11.1 ij	13.0 d-k	0.6 hi	4.3 d-h
N15044oIF	2.0 g-j	1.2 b-g	85.1 cd	5.9 lm	13.6 c-e	0.9 de	4.2 e-h
N15053	2.1 c-i	1.2 b-f	81.3 fg	10.2 jk	13.1 d-i	0.6 gh	4.4 b-g
N15060	2.3 a	1.5 a	79.5 g-i	15.3 d-g	12.8 f-l	0.4 ij	4.8 a
N16005	2.0 f-j	1.0 g	82.9 ef	7.7 kl	14.0 bc	0.7 fg	4.2 e-h
N16012	2.1 b-h	1.2 b-f	78.5 ij	18.7 bc	13.0 d-k	0.3 jk	4.5 a-e
N16021	2.2 a-f	1.3 b-f	82.9 ef	7.8 kl	13.5 c-f	0.7 fg	4.5 a-e
N17036	2.2 a-f	1.3 b-f	78.9 ij	17.4 b-f	12.9 e-l	0.4 jk	4.5 a-e
N17037	2.1 b-g	1.2 b-f	78.6 ij	17.7 b-e	13.1 d-j	0.4 jk	4.5 a-e
N17040	2.0 f-j	1.2 b-g	79.5 g-i	16.8 c-f	12.3 j-l	0.4 i-k	4.3 d-h
N17041	2.0 f-j	1.3 b-e	79.3 hi	17.7 b-e	12.3 i-l	0.4 jk	4.3 c-h
N17044	2.2 a-d	1.2 d-g	77.2 j	26.8 a	13.3 c-g	0.2 k	4.7 a-c
N17045	1.9 ij	1.2 d-g	80.0 g-i	15.0 e-g	12.4 h-l	0.4 h-j	4.1 gh
N17047	2.1 d-i	1.3 a-d	79.1 i	18.0 b-d	12.5 h-l	0.4 jk	4.4 b-g
Mean	2.1	1.2	81.5	13.4	13.1	0.6	4.4
LSD²	0.2	0.2	1.8	2.8	0.8	0.2	0.4

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 26. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Rocky Mount, NC, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.3 a	2.4 c-e	53.2 k	29.2 a	1.1 bc	1.3 j
Bailey II	5.8 f-i	2.3 c-g	80.9 a-d	5.6 g-l	1.0 c-f	1.6 d-h
Emery	5.8 f-i	2.4 c-f	80.7 a-d	5.2 h-l	1.1 b-d	1.7 c-h
Sullivan	6.2 e-h	2.3 c-f	77.6 c-g	7.7 e-i	1.1 b-d	1.8 a-e
Walton	5.8 f-i	2.5 bc	78.5 b-f	6.5 g-k	1.2 b	2.0 a
Wynne	6.0 e-i	2.3 c-f	80.0 a-d	6.1 g-l	1.1 b-d	1.6 d-h
N14001	5.8 g-i	2.3 c-f	82.0 ab	4.4 kl	1.1 b-f	1.6 e-i
N14002oIJ	6.5 b-g	2.5 b-d	77.3 d-h	7.9 e-h	1.1 b-d	1.6 d-h
N14004oIJ	6.5 b-f	2.6 bc	75.9 e-i	9.3 d-f	1.1 bc	1.6 e-i
N14007	6.9 b-d	2.0 g	71.9 j	13.9 b	0.9 d-f	1.5 h-j
N14009	5.4 ij	2.2 d-g	81.8 ab	4.4 kl	1.1 b-f	1.9 a-c
N15066	6.9 bc	2.4 c-e	74.4 g-j	10.0 c-e	1.1 b-e	1.8 a-d
N14023oI	5.9 e-i	2.4 c-f	81.0 a-c	5.0 j-l	1.0 b-f	1.8 b-f
N14027oIJ	7.1 b	2.2 e-g	72.5 ij	12.5 bc	1.0 b-f	1.5 g-i
N15017oI	6.1 e-i	2.1 e-g	79.9 a-d	5.6 g-l	1.0 b-f	1.9 ab
N15039oI	6.4 b-g	2.5 bc	75.2 f-j	9.9 c-e	1.1 bc	1.6 e-i
N15041oI	6.3 c-h	2.1 e-g	79.5 a-e	6.6 g-k	1.0 c-f	1.6 f-i
N15044oIF	6.6 b-e	2.2 e-g	75.9 e-i	9.6 d-f	1.0 b-f	1.6 c-h
N15053	5.8 f-i	2.3 c-g	80.8 a-d	5.3 h-l	1.0 b-f	1.7 b-h
N15060	5.5 ij	2.2 e-g	81.6 ab	5.2 i-l	0.9 ef	1.7 b-g
N16005	6.9 bc	2.7 b	73.9 h-j	11.1 cd	1.2 b	1.3 j
N16012	5.8 g-i	2.3 c-g	81.3 a-c	4.8 j-l	1.1 b-e	1.7 b-g
N16021	6.2 c-h	2.1 e-g	77.3 d-h	8.0 e-g	1.1 b-f	1.8 a-e
N17036	5.9 e-i	2.6 bc	80.7 a-d	4.8 j-l	1.1 bc	1.7 b-h
N17037	5.0 j	2.1 fg	83.0 a	5.1 i-l	0.9 f	1.4 ij
N17040	5.8 g-i	2.4 c-e	80.8 a-d	5.1 i-l	1.1 b-f	1.8 a-f
N17041	5.9 e-i	2.3 c-g	79.7 a-d	6.1 g-l	1.1 b-f	1.7 b-g
N17044	5.6 h-j	3.2 a	81.1 a-c	3.6 l	1.4 a	1.7 b-g
N17045	6.2 d-h	2.3 c-g	78.4 b-f	7.2 f-j	1.1 b-f	1.7 b-h
N17047	5.9 f-i	2.3 c-g	80.8 a-d	5.3 h-l	1.0 c-f	1.8 a-e
Mean	6.2	2.3	77.9	7.7	1.1	1.7
LSD²	0.7	0.3	3.7	2.7	0.2	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 26. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Rocky Mount, NC, 2020¹ (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.4 a	1.1 a-d	97.3 a	1.8 l	16.3 a	1.8 a	4.6 a-c
Bailey II	1.9 e-g	1.0 b-e	80.6 f-j	14.5 b-f	11.8 d-f	0.5 g-k	3.8 fg
Emery	2.1 a-f	1.0 a-e	79.8 h-j	15.5 b-e	12.4 b-e	0.4 h-l	4.2 a-f
Sullivan	2.2 a-e	1.1 a-d	81.5 e-i	10.2 f-k	12.9 b-e	0.6 e-h	4.4 a-f
Walton	2.3 a-c	1.2 ab	80.3 h-j	12.1 d-i	13.0 b-e	0.5 g-k	4.7 ab
Wynne	1.9 e-g	1.0 a-e	80.6 f-j	13.2 d-g	12.3 b-e	0.5 g-k	4.0 c-g
N14001	1.8 fg	1.0 a-e	79.4 i-k	19.0 ab	12.0 c-f	0.4 kl	3.9 d-g
N14002olJ	2.1 a-g	1.1 a-e	81.4 e-i	11.4 e-j	13.2 b-d	0.6 e-i	4.3 a-f
N14004olJ	2.0 a-g	1.0 a-e	82.6 d-g	8.3 g-k	13.2 b-d	0.7 d-f	4.2 a-f
N14007	1.9 d-g	0.9 c-e	87.1 b	5.2 kl	12.7 b-e	1.1 b	3.8 e-g
N14009	2.1 a-f	1.1 a-d	79.5 i-k	18.6 a-c	11.9 d-f	0.4 kl	4.3 a-f
N15066	2.3 a-d	1.2 a-c	82.7 d-f	7.5 i-k	13.8 b	0.7 de	4.5 a-e
N14023ol	1.9 e-g	1.1 a-d	79.7 h-j	16.7 b-d	12.2 b-e	0.4 h-l	4.0 c-g
N14027olJ	2.0 d-g	1.2 a	85.2 bc	5.9 kl	13.5 bc	0.9 bc	4.3 a-f
N15017ol	2.2 a-e	1.2 a-c	80.0 h-j	14.3 b-f	12.5 b-e	0.4 g-l	4.4 a-f
N15039ol	2.1 a-g	1.1 a-d	83.2 c-e	7.6 h-k	13.2 b-d	0.7 c-e	4.3 a-f
N15041ol	1.8 e-g	1.1 a-d	81.0 e-j	12.7 d-h	12.3 b-e	0.5 f-k	3.9 d-g
N15044olF	2.0 c-g	1.1 a-d	83.2 c-e	7.9 h-k	12.9 b-e	0.7 c-e	4.1 b-g
N15053	1.9 d-g	1.1 a-d	80.0 h-j	15.4 b-e	12.2 b-e	0.4 g-l	4.1 b-g
N15060	2.0 a-g	1.0 b-e	80.5 g-j	15.8 b-e	11.5 ef	0.5 g-l	3.9 d-g
N16005	2.0 a-g	0.9 de	83.7 cd	6.7 j-l	13.8 b	0.8 cd	4.1 a-g
N16012	1.9 d-g	1.1 a-e	79.7 h-j	16.8 b-d	12.1 c-e	0.4 i-l	4.1 b-g
N16021	2.3 a-c	1.1 a-d	81.7 d-h	9.9 f-k	12.9 b-e	0.6 d-g	4.5 a-d
N17036	2.1 a-f	1.0 a-e	79.1 jk	16.7 b-d	12.8 b-e	0.4 j-l	4.3 a-f
N17037	1.8 g	0.8 e	81.4 e-i	16.4 b-e	10.5 f	0.5 g-k	3.5 g
N17040	2.0 b-g	1.1 a-d	79.8 h-j	16.0 b-e	12.3 b-e	0.4 h-l	4.2 a-f
N17041	2.1 a-f	1.1 a-d	80.4 g-j	13.8 c-f	12.5 b-e	0.5 g-k	4.2 a-f
N17044	2.3 ab	1.1 a-d	77.4 k	22.3 a	13.6 bc	0.3 l	4.8 a
N17045	2.1 a-f	1.1 a-d	81.3 e-j	12.2 d-i	12.7 b-e	0.6 e-j	4.2 a-f
N17047	1.9 d-g	1.0 a-e	80.1 h-j	15.3 b-e	12.1 c-e	0.4 g-l	4.0 c-g
Mean	2.1	1.1	81.7	12.7	12.7	0.6	4.2
LSD²	0.4	0.3	2.2	5.1	1.6	0.2	0.7

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 27. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Bladen County, NC, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.4 a	2.2 e-j	51.8 k	30.7 a	1.1 b-d	1.2 l
Bailey II	5.8 h-j	2.1 g-k	81.3 a-e	5.3 g-m	1.0 c-e	1.6 e-i
Emery	5.9 e-j	2.3 d-h	80.7 a-f	5.1 h-m	1.1 b-d	1.5 e-j
Sullivan	6.3 d-h	2.1 g-k	77.1 f-h	8.3 e-g	1.0 cd	1.7 b-h
Walton	5.9 f-j	2.4 b-f	78.9 b-g	6.4 f-l	1.1 bc	1.8 a-c
Wynne	6.2 d-h	2.4 b-f	77.9 d-h	7.3 f-k	1.1 bc	1.6 d-i
N14001	5.8 h-j	2.3 c-h	81.9 a-c	4.3 k-m	1.1 b-d	1.5 g-k
N14002oIJ	5.9 e-i	2.3 c-g	81.7 a-d	4.1 lm	1.1 b-d	1.6 e-i
N14004oIJ	6.4 de	2.5 b-d	77.1 f-h	8.1 e-h	1.1 bc	1.5 e-j
N14007	8.1 b	2.1 g-k	62.9 j	20.9 b	1.0 cd	1.4 j-l
N14009	5.4 ij	2.1 h-k	83.0 a	3.7 lm	1.0 c-e	1.6 b-h
N15066	6.3 d-h	2.3 d-h	78.5 c-g	6.1 g-m	1.1 b-d	1.8 ab
N14023ol	5.9 e-j	2.0 k	81.7 a-d	4.5 j-m	1.0 de	1.8 a-c
N14027oIJ	7.1 c	2.1 g-k	71.7 i	13.2 c	1.0 c-e	1.5 f-j
N15017ol	6.3 d-h	2.2 e-j	78.6 b-g	6.9 f-l	1.0 cd	1.6 c-i
N15039ol	6.3 d-g	2.3 d-i	76.1 gh	9.5 d-f	1.1 b-d	1.5 g-k
N15041ol	6.4 d-f	2.0 i-k	77.6 e-h	7.9 e-i	1.0 c-e	1.8 a-d
N15044oIF	6.4 d-f	2.0 jk	78.3 c-g	7.6 f-j	1.0 de	1.6 c-i
N15053	6.1 d-h	2.2 d-j	78.4 c-g	7.3 f-k	1.1 b-d	1.6 e-i
N15060	5.8 g-j	2.2 f-j	79.2 b-g	6.0 g-m	1.0 cd	1.9 a
N16005	7.1 c	2.5 bc	72.1 i	12.2 cd	1.2 b	1.3 kl
N16012	5.8 h-j	2.2 f-k	80.6 a-f	5.1 h-m	1.1 b-d	1.7 a-e
N16021	6.7 cd	2.1 h-k	74.4 hi	10.8 c-e	1.0 cd	1.6 d-i
N17036	5.3 j	2.4 b-e	82.3 ab	4.4 k-m	1.0 c-e	1.4 i-k
N17037	5.8 g-j	2.6 b	81.2 a-e	4.1 lm	1.2 b	1.6 e-i
N17040	5.7 h-j	2.3 d-h	80.6 a-f	5.2 g-m	1.1 b-d	1.6 b-h
N17041	5.5 ij	2.1 g-k	82.0 a-c	4.2 k-m	1.0 c-e	1.7 b-g
N17044	5.4 ij	3.1 a	82.3 ab	3.0 m	1.3 a	1.5 h-k
N17045	5.8 f-j	2.0 k	81.4 a-d	5.1 h-m	0.9 e	1.6 b-h
N17047	5.9 e-i	2.2 f-k	81.0 a-e	4.8 i-m	1.0 c-e	1.7 b-f
Mean	6.2	2.3	77.8	7.7	1.1	1.6
LSD²	0.6	0.2	3.8	3.2	0.1	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 27. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Bladen County, NC, 2020¹ (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.2 a-c	1.4 a-c	98.7 a	1.7 n	16.3 a	1.9 a	4.7 a-e
Bailey II	1.9 f-i	1.0 d	80.3 g-l	15.7 c-j	11.9 i-k	0.4 g-l	4.0 f
Emery	2.1 b-i	1.3 b-d	79.5 g-m	16.1 b-i	12.6 d-k	0.4 h-m	4.4 a-f
Sullivan	2.1 a-h	1.4 a-c	82.0 d-g	9.6 i-m	13.0 d-i	0.6 d-g	4.6 a-f
Walton	2.2 a-e	1.4 a-c	80.4 g-l	13.2 f-l	12.9 d-j	0.5 g-l	4.7 a-e
Wynne	2.1 a-g	1.4 bc	80.9 f-k	10.8 g-m	13.2 d-g	0.6 f-k	4.6 a-f
N14001	1.9 g-i	1.1 b-d	79.2 i-m	18.9 b-f	12.2 f-k	0.4 j-m	4.1 c-f
N14002oIJ	2.1 b-i	1.3 b-d	78.5 k-m	20.2 bc	12.7 d-j	0.3 lm	4.5 a-f
N14004oIJ	2.1 b-i	1.2 b-d	81.6 e-i	9.6 j-m	13.3 d-f	0.6 e-h	4.4 b-f
N14007	2.2 a-e	1.4 a-c	91.4 b	3.0 n	14.8 b	1.4 b	4.6 a-f
N14009	2.0 d-i	1.2 b-d	79.2 i-m	22.3 ab	11.6 k	0.3 lm	4.1 c-f
N15066	2.3 ab	1.7 a	79.5 h-m	13.3 e-l	13.6 cd	0.4 g-l	5.0 a
N14023oI	1.9 f-i	1.3 b-d	79.4 h-m	18.4 b-f	12.1 h-k	0.4 j-m	4.2 b-f
N14027oIJ	2.1 c-i	1.3 b-d	85.7 c	5.4 mn	13.6 c-e	1.0 c	4.4 b-f
N15017oI	2.0 c-i	1.4 a-c	80.8 f-k	14.3 c-k	12.9 d-j	0.5 f-l	4.4 a-f
N15039oI	2.0 c-i	1.2 b-d	83.0 d-f	8.1 k-n	12.9 d-j	0.7 d-f	4.3 b-f
N15041oI	2.0 c-i	1.3 b-d	81.8 d-h	9.9 h-m	12.8 d-j	0.6 e-h	4.3 b-f
N15044oIF	1.8 i	1.3 b-d	81.7 d-h	11.6 g-m	12.5 d-k	0.6 e-i	4.1 d-f
N15053	2.0 c-i	1.2 b-d	81.3 f-j	10.9 g-m	12.7 d-k	0.6 f-j	4.3 b-f
N15060	2.4 a	1.4 ab	80.1 g-l	13.5 d-l	12.8 d-j	0.5 g-l	4.8 ab
N16005	2.2 a-c	1.4 a-c	84.1 cd	6.0 mn	14.4 bc	0.8 cd	4.8 a-c
N16012	2.2 a-f	1.4 a-c	79.5 g-m	15.8 b-j	12.6 d-k	0.4 h-m	4.6 a-e
N16021	2.1 a-g	1.3 b-d	83.9 c-e	7.2 l-n	13.2 d-g	0.8 c-e	4.5 a-f
N17036	2.0 e-i	1.1 cd	79.6 g-l	18.6 b-f	11.9 jk	0.4 j-m	4.1 ef
N17037	2.1 a-h	1.4 ab	78.2 lm	19.9 b-d	13.1 d-h	0.3 lm	4.7 a-d
N17040	2.0 c-i	1.4 a-c	79.7 g-l	16.4 b-g	12.5 e-k	0.4 h-m	4.5 a-f
N17041	2.1 c-i	1.4 ab	79.1 j-m	19.7 b-e	12.1 g-k	0.3 k-m	4.5 a-f
N17044	2.2 a-d	1.2 b-d	77.1 m	28.2 a	13.3 d-f	0.2 m	4.8 ab
N17045	1.9 hi	1.3 b-d	80.1 g-l	16.3 b-h	11.9 i-k	0.4 g-m	4.1 ef
N17047	2.0 c-i	1.4 a-c	79.4 h-m	16.8 b-g	12.5 f-k	0.4 i-m	4.4 b-f
Mean	2.1	1.3	81.5	13.7	12.9	0.6	4.5
LSD²	0.3	0.3	2.5	6.5	1.1	0.2	0.6

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 28. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Blackville, SC, 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	7.8 a	2.0 o	59.4 f	27.2 a	0.8 l	0.8 m
Bailey II	7.3 a-c	2.4 g-l	81.0 ab	4.0 fg	1.2 d-h	1.6 b-f
Emery	6.1 hi	2.6 c-e	78.2 a-c	6.6 e-g	1.2 cd	1.4 i-k
Sullivan	6.4 e-i	2.3 i-n	80.8 ab	4.3 fg	1.2 d-g	1.7 a-c
Walton	6.1 hi	2.5 e-j	79.4 ab	5.0 fg	1.2 c-e	1.8 a
Wynne	6.0 i	2.4 f-j	76.9 a-c	7.6 d-f	1.2 e-i	1.5 f-k
N14001	6.8 c-g	2.5 d-j	80.7 ab	4.7 fg	1.2 d-h	1.5 g-k
N14002oIJ	6.1 f-i	2.5 d-h	79.8 ab	4.8 fg	1.2 c-e	1.6 c-i
N14004oIJ	6.4 f-i	2.7 bc	79.1 ab	5.4 fg	1.3 c	1.5 h-k
N14007	6.5 d-i	2.4 h-m	68.3 e	15.4 b	1.2 d-h	1.4 kl
N14009	6.1 g-i	2.4 f-k	80.5 ab	4.8 fg	1.1 f-j	1.5 e-j
N15066	6.1 f-i	2.5 d-i	79.4 ab	5.2 fg	1.2 c-e	1.8 ab
N14023ol	6.3 f-i	2.2 mn	81.2 ab	4.4 fg	1.1 k	1.6 c-f
N14027oIJ	7.5 a-c	2.3 i-n	70.6 de	13.4 bc	1.1 f-j	1.4 jk
N15017ol	6.4 f-i	2.4 h-n	79.5 ab	5.4 fg	1.2 e-i	1.6 c-h
N15039ol	6.4 e-i	2.4 e-j	77.5 a-c	7.1 d-g	1.2 c-e	1.6 c-h
N15041ol	6.8 c-h	2.2 l-n	77.4 a-c	7.4 d-g	1.1 jk	1.6 d-j
N15044oIF	7.2 a-e	2.2 n	73.2 c-e	11.1 b-d	1.1 i-k	1.7 b-d
N15053	7.6 ab	2.5 d-j	69.8 de	13.9 bc	1.2 c-e	1.4 i-k
N15060	6.2 f-i	2.3 i-n	79.0 ab	5.5 fg	1.1 g-k	1.7 a-c
N16005	6.9 b-f	2.8 ab	76.3 bc	7.9 d-f	1.3 b	1.2 l
N16012	6.1 f-i	2.3 i-n	80.0 ab	4.8 fg	1.2 d-h	1.7 b-e
N16021	7.2 a-d	2.4 h-n	73.5 cd	10.6 c-e	1.2 e-i	1.5 d-j
N17036	6.2 f-i	2.6 c-f	80.9 ab	3.9 fg	1.3 c	1.6 d-j
N17037	6.5 d-i	2.7 b-d	79.4 ab	5.7 fg	1.2 d-f	1.4 i-k
N17040	6.1 f-i	2.2 k-n	80.1 ab	5.3 fg	1.1 i-k	1.6 c-f
N17041	6.0 i	2.3 j-n	80.9 ab	4.8 fg	1.1 h-k	1.6 c-h
N17044	6.0 i	2.9 a	81.1 ab	3.2 g	1.4 a	1.6 c-g
N17045	6.3 f-i	2.3 i-n	80.0 ab	5.1 fg	1.1 g-k	1.6 d-j
N17047	6.3 f-i	2.2 l-n	81.1 ab	4.3 fg	1.1 jk	1.6 b-f
N14017	5.9 i	2.6 c-g	82.0 a	3.5 fg	1.2 c-e	1.5 e-k
Mean	6.5	2.4	77.6	7.2	1.2	1.5
LSD²	0.7	0.2	5.2	4.4	0.1	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 28. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated from Blackville, SC, 2020¹, (cont.).

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	1.7 i	0.8 d	98.8 a	2.2 h	12.6 i	2.2 a	3.3 h
Bailey II	2.3 a-h	1.4 ab	77.9 fg	20.5 a-c	13.3 e-i	0.3 fg	4.9 a-f
Emery	2.1 gh	1.3 a-c	79.9 ef	11.8 d-f	13.7 c-g	0.5 e-g	4.7 c-g
Sullivan	2.3 a-g	1.3 a-c	78.3 fg	18.7 bc	13.2 e-i	0.3 fg	4.8 a-g
Walton	2.5 a	1.5 a	78.5 fg	15.8 c-e	13.7 c-g	0.4 fg	5.2 a
Wynne	2.3 a-h	1.3 a-c	80.5 d-f	10.2 e-g	14.0 a-e	0.5 d-f	4.7 a-g
N14001	2.1 gh	1.3 a-c	78.7 fg	17.3 b-d	13.2 e-i	0.4 fg	4.6 d-g
N14002oIJ	2.3 a-g	1.4 ab	78.1 fg	16.7 c-e	13.9 c-f	0.3 fg	5.0 a-d
N14004oIJ	2.2 d-h	1.3 a-c	78.6 fg	15.3 c-e	14.0 b-e	0.4 fg	4.8 a-g
N14007	2.3 a-g	1.3 a-c	86.5 b	4.4 gh	15.0 a	1.0 b	4.8 a-g
N14009	2.2 e-h	1.4 ab	78.7 fg	16.7 c-e	13.2 e-i	0.4 fg	4.7 b-g
N15066	2.5 a-d	1.3 a-c	78.7 fg	15.7 c-e	13.6 c-h	0.4 fg	5.0 a-d
N14023oI	2.1 h	1.2 a-c	78.7 fg	19.0 a-c	12.8 g-i	0.3 fg	4.3 fg
N14027oIJ	2.2 f-h	1.4 ab	85.1 bc	5.3 f-h	14.5 a-c	0.9 bc	4.7 a-g
N15017oI	2.3 a-h	1.4 ab	78.9 fg	15.4 c-e	13.5 d-i	0.4 fg	4.8 a-g
N15039oI	2.3 a-g	1.3 a-c	80.3 d-f	11.3 d-f	13.8 c-g	0.5 d-f	4.9 a-e
N15041oI	2.1 gh	1.3 a-c	80.7 d-f	10.4 e-g	13.6 d-i	0.5 d-f	4.6 d-g
N15044oIF	2.3 a-h	1.3 a-c	83.5 b-d	6.6 f-h	14.0 a-e	0.8 b-d	4.7 a-g
N15053	2.4 a-f	1.3 a-c	85.2 bc	7.0 f-h	14.9 ab	0.9 bc	4.9 a-f
N15060	2.5 ab	1.6 a	78.9 fg	14.4 c-e	13.7 c-g	0.4 fg	5.2 ab
N16005	2.3 b-h	1.3 a-c	80.3 d-f	10.3 e-g	14.5 a-c	0.5 d-f	4.9 a-f
N16012	2.4 a-e	1.4 ab	78.5 fg	16.6 c-e	13.5 d-i	0.4 fg	5.0 a-d
N16021	2.4 a-f	1.3 a-c	82.8 c-e	7.3 f-h	14.4 a-d	0.7 c-e	4.8 a-g
N17036	2.3 a-g	1.2 a-c	77.6 fg	21.1 a-c	13.7 c-h	0.3 fg	4.8 a-g
N17037	2.1 gh	1.0 cd	79.2 fg	16.1 c-e	13.5 d-i	0.4 fg	4.3 g
N17040	2.2 e-h	1.3 a-c	79.3 fg	15.2 c-e	13.0 f-i	0.4 fg	4.6 d-g
N17041	2.2 e-h	1.1 b-d	79.1 fg	17.6 b-d	12.7 hi	0.4 fg	4.4 e-g
N17044	2.5 a-c	1.3 a-c	76.6 g	25.6 a	14.1 a-e	0.2 g	5.1 a-c
N17045	2.3 c-h	1.3 a-c	78.8 fg	16.6 c-e	13.3 e-i	0.4 fg	4.7 a-g
N17047	2.1 gh	1.3 a-c	78.4 fg	19.2 a-c	13.0 f-i	0.3 fg	4.5 d-g
N14017	2.2 e-h	1.1 b-d	77.8 fg	23.7 ab	13.0 f-i	0.3 fg	4.5 d-g
Mean	2.3	1.3	80.4	14.3	13.6	0.5	4.7
LSD²	0.2	0.4	3.3	6.8	1.0	0.3	0.5

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 29. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated averaged across all locations, 2020.¹

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	8.9 a	2.2 g-k	54.9 k	28.2 a	1.1 e-k	1.2 l
Bailey II	5.9 l-p	2.3 e-i	80.5 a-c	5.3 j-m	1.1 e-k	1.6 d-h
Emery	5.9 l-p	2.4 c-e	80.5 a-c	5.2 k-m	1.1 cd	1.6 g-j
Sullivan	6.1 i-n	2.2 h-k	79.2 a-e	6.3 h-l	1.1 d-j	1.7 b-f
Walton	5.8 n-p	2.3 d-h	79.3 a-e	6.0 i-l	1.1 c-h	1.9 a
Wynne	6.3 f-j	2.4 c-e	78.1 d-f	7.1 g-j	1.1 bc	1.6 g-j
N14001	6.0 j-o	2.4 c-f	80.8 a-c	5.1 k-m	1.1 c-g	1.5 i-k
N14002oIJ	6.3 f-k	2.4 c-f	79.3 a-e	5.9 i-l	1.1 c-f	1.6 e-i
N14004oIJ	6.5 e-g	2.3 c-g	75.7 gh	9.6 d-f	1.1 c-i	1.5 jk
N14007	7.6 b	2.2 jk	67.9 j	16.5 b	1.1 f-k	1.5 k
N14009	5.6 p	2.2 h-k	81.4 a	4.6 lm	1.1 f-k	1.7 b-d
N15066	6.6 e-g	2.2 h-k	77.3 e-g	7.5 g-i	1.0 g-k	1.8 ab
N14023oI	6.0 j-n	2.1 jk	81.3 ab	4.6 lm	1.0 jk	1.7 b-d
N14027oIJ	7.2 c	2.2 jk	71.4 i	13.3 c	1.0 f-k	1.5 i-k
N15017oI	6.3 g-l	2.2 h-k	79.1 b-f	6.3 h-l	1.1 d-j	1.7 b-f
N15039oI	6.5 e-h	2.4 c-f	75.6 gh	9.4 ef	1.1 c-e	1.6 h-k
N15041oI	6.6 ef	2.1 k	77.3 e-g	8.1 f-h	1.0 k	1.6 d-i
N15044oIF	7.0 cd	2.1 k	73.6 hi	11.4 d	1.0 h-k	1.6 e-i
N15053	6.4 f-i	2.2 f-j	77.0 fg	8.2 fg	1.1 c-i	1.7 d-h
N15060	5.9 l-p	2.1 jk	79.4 a-e	5.9 i-l	1.0 jk	1.9 a
N16005	7.0 cd	2.6 b	73.7 h	10.9 de	1.2 b	1.3 l
N16012	5.9 l-p	2.3 e-h	80.6 a-c	5.0 k-m	1.1 c-g	1.7 c-g
N16021	6.8 de	2.2 h-k	74.2 h	10.6 de	1.1 c-i	1.6 d-h
N17036	6.2 h-m	2.4 cd	78.6 c-f	6.7 g-k	1.1 cd	1.6 f-i
N17037	5.9 l-p	2.5 c	80.3 a-c	5.3 j-m	1.1 cd	1.6 g-j
N17040	5.9 l-p	2.2 h-k	80.2 a-d	5.6 j-l	1.0 f-k	1.7 b-e
N17041	5.9 m-p	2.2 jk	80.6 a-c	5.3 j-m	1.0 i-k	1.7 b-e
N17044	5.7 op	3.0 a	81.4 a	3.5 m	1.3 a	1.6 g-j
N17045	6.1 i-n	2.1 jk	79.8 a-d	5.9 i-l	1.0 jk	1.7 b-f
N17047	6.0 k-p	2.2 i-k	81.0 ab	4.8 k-m	1.0 i-k	1.8 bc
Mean	6.4	2.3	77.3	7.9	1.1	1.6
LSD²	0.3	0.1	2.2	1.9	0.1	0.1

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 29. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated averaged across all locations, 2020¹. (cont.)

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.3 a-d	1.2 e-g	97.1 a	2.0 r	15.6 a	1.8 a	4.5 b-f
Bailey II	2.0 h-k	1.2 d-g	79.8 ij	15.6 c-i	12.5 l-n	0.4 kl	4.3 e-h
Emery	2.1 f-j	1.2 c-g	79.4 j	16.1 b-h	12.8 h-n	0.4 l	4.4 d-h
Sullivan	2.1 e-i	1.2 b-g	80.4 h-j	13.4 i-k	12.8 i-n	0.5 i-l	4.4 d-h
Walton	2.3 a-c	1.3 ab	80.0 ij	13.8 g-j	12.9 g-m	0.5 j-l	4.7 a-c
Wynne	2.2 d-i	1.2 c-g	80.7 h-j	11.6 j-l	13.2 d-i	0.5 h-k	4.5 b-f
N14001	2.0 jk	1.1 g	79.6 ij	17.3 b-d	12.6 k-n	0.4 l	4.2 h
N14002oIJ	2.1 e-i	1.2 b-f	79.7 ij	14.7 e-i	13.2 e-j	0.4 j-l	4.5 c-f
N14004oIJ	2.1 g-k	1.2 e-g	82.9 d-g	9.5 l-o	13.2 d-i	0.7 d-f	4.3 e-h
N14007	2.2 c-h	1.2 c-g	88.1 b	4.8 q	14.2 b	1.1 b	4.5 d-h
N14009	2.1 e-i	1.2 c-g	79.4 j	18.2 b	12.2 n	0.4 lm	4.4 d-h
N15066	2.3 a-f	1.3 a-c	80.9 hi	11.0 k-m	13.4 d-h	0.6 h-j	4.6 b-d
N14023oI	2.0 k	1.2 b-g	79.3 j	17.9 bc	12.4 mn	0.4 lm	4.2 gh
N14027oIJ	2.1 g-k	1.2 b-e	85.7 c	6.0 pq	13.8 b-d	1.0 c	4.4 d-h
N15017oI	2.1 e-i	1.3 a-e	80.2 ij	13.6 h-j	12.9 g-m	0.5 j-l	4.5 d-g
N15039oI	2.2 d-i	1.2 c-g	82.6 e-g	8.8 m-o	13.4 d-h	0.7 e-g	4.5 c-f
N15041oI	2.0 jk	1.3 a-d	81.8 f-h	10.1 l-n	13.0 f-l	0.6 f-h	4.3 f-h
N15044oIF	2.0 i-k	1.2 b-g	84.3 cd	7.2 o-q	13.4 d-g	0.8 cd	4.3 e-h
N15053	2.2 d-i	1.2 b-e	81.8 gh	10.5 lm	13.1 e-k	0.6 g-i	4.5 c-f
N15060	2.4 a	1.4 a	80.0 ij	14.1 f-i	12.8 h-n	0.5 j-l	4.8 ab
N16005	2.1 f-j	1.1 fg	83.3 d-f	7.2 o-q	14.1 bc	0.8 de	4.4 d-h
N16012	2.2 d-i	1.2 b-g	79.3 j	16.7 b-e	12.7 i-n	0.4 l	4.5 b-f
N16021	2.3 a-e	1.3 b-e	83.5 de	7.8 n-p	13.6 c-e	0.8 de	4.6 b-d
N17036	2.2 b-g	1.2 d-g	80.4 h-j	16.1 b-h	13.1 e-j	0.5 j-l	4.5 b-e
N17037	2.1 d-i	1.2 e-g	79.5 ij	16.3 b-f	12.8 g-m	0.4 kl	4.4 d-h
N17040	2.1 g-k	1.2 b-g	80.0 ij	15.2 d-i	12.5 l-n	0.4 j-l	4.4 d-h
N17041	2.1 g-k	1.2 b-g	79.8 ij	16.2 b-g	12.4 l-n	0.4 kl	4.4 d-h
N17044	2.3 ab	1.2 c-g	77.3 k	24.1 a	13.5 d-f	0.3 m	4.9 a
N17045	2.1 h-k	1.2 c-g	80.2 ij	14.4 e-i	12.6 j-n	0.5 j-l	4.3 e-h
N17047	2.1 h-k	1.3 b-e	79.3 j	17.2 b-d	12.5 l-n	0.4 lm	4.3 e-h
Mean	2.1	1.2	81.5	12.9	13.1	0.6	4.5
LSD	0.1	0.1	1.5	2.5	0.6	0.1	0.3

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 30. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Two-year averages across all locations, (2019 – 2020)¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	8.9 a	2.4 cd	55.4 g	27.0 a	1.2 b-d	1.3 g
Bailey II	6.4 e-g	2.4 b-d	75.8 e	9.1 bc	1.1 b-d	1.6 d-f
Emery	5.8 ij	2.5 b	80.4 ab	4.8 fg	1.2 bc	1.6 d-f
Sullivan	6.0 hi	2.3 de	78.8 bc	6.3 ef	1.1 cd	1.7 bc
Walton	5.7 j	2.5 bc	79.3 a-c	5.3 fg	1.2 b	1.9 a
Wynne	6.2 f-h	2.5 bc	78.0 c	6.9 e	1.2 b-d	1.6 f
N14002oIJ	6.6 c-e	2.5 bc	75.7 ef	8.8 c	1.2 b-d	1.6 f
N14004oIJ	6.4 d-f	2.5 bc	76.1 e	8.6 cd	1.2 b-d	1.6 f
N14023ol	6.0 hi	2.2 f	80.9 a	4.6 g	1.1 e	1.8 bc
N14027oIJ	6.7 cd	2.2 f	75.5 ef	9.4 bc	1.1 e	1.6 d-f
N15017ol	6.2 gh	2.3 ef	78.8 bc	6.2 ef	1.1 d	1.8 b
N15039ol	6.1 h	2.4 bc	78.0 cd	7.1 de	1.2 b-d	1.6 ef
N15041ol	6.6 c-e	2.2 f	76.3 de	8.6 cd	1.1 e	1.7 c-e
N15044oIF	6.7 bc	2.2 f	75.1 ef	9.7 bc	1.1 e	1.7 cd
N16005	6.9 b	2.8 a	74.0 f	10.2 b	1.3 a	1.3 g
Mean	6.5	2.4	75.9	8.8	1.1	1.6
LSD²	0.3	0.1	1.7	1.5	0.1	0.1

¹ Refer to page 3 for an explanation of the computations of these characters.

² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 30. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Two-year averages across all locations, (2019 – 2020)¹, (cont.)

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.5 ab	1.3 cd	95.4 a	2.1 g	16.3 a	1.7 a	5.0 b
Bailey II	2.3 c-e	1.3 cd	82.2 b	10.9 cd	13.5 c-e	0.6 b	4.7 b-d
Emery	2.3 c-e	1.3 cd	78.8 e	17.2 a	13.1 ef	0.4 g	4.8 b-d
Sullivan	2.3 c-e	1.4 ab	80.0 de	13.3 b	13.2 e	0.5 ef	4.9 bc
Walton	2.6 a	1.5 a	78.9 e	16.4 a	13.5 de	0.4 fg	5.3 a
Wynne	2.4 cd	1.3 cd	80.3 d	12.2 bc	13.5 c-e	0.5 de	4.8 b-d
N14002oIJ	2.4 bc	1.3 cd	81.5 bc	10.8 c-e	14.0 bc	0.6 b-d	4.9 b-d
N14004oIJ	2.3 c-e	1.3 bc	81.5 bc	10.1 c-e	13.8 cd	0.6 bc	4.8 b-d
N14023oI	2.2 e	1.3 a-c	78.9 e	18.2 a	12.7 f	0.4 g	4.6 d
N14027oIJ	2.2 de	1.3 bc	82.6 b	10.6 c-e	13.5 de	0.7 b	4.6 d
N15017oI	2.4 cd	1.4 a-c	79.8 de	14.2 b	13.3 e	0.5 e-g	4.9 b-d
N15039oI	2.3 c-e	1.3 cd	80.7 cd	13.3 b	13.3 e	0.5 c-e	4.7 b-d
N15041oI	2.2 de	1.4 a-c	81.8 bc	9.4 d-f	13.4 de	0.6 bc	4.7 cd
N15044oIF	2.3 c-e	1.3 bc	82.7 b	8.6 ef	13.5 c-e	0.7 b	4.7 cd
N16005	2.3 c-e	1.2 d	82.4 b	7.7 f	14.4 b	0.7 b	4.7 b-d
Mean	2.3	1.3	81.8	11.7	13.7	0.6	4.8
LSD²	0.1	0.1	1.2	2.1	0.4	0.1	0.3

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.³ Lower iodine value indicates longer shelf life.⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 31. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Three-year averages across all locations, (2018 – 2020)¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
Bailey	9.0 a	2.5 bc	55.3 e	26.9 a	1.2 a	1.3 g
Bailey II	6.2 c-e	2.5 bc	77.7 d	7.3 b	1.2 ab	1.6 ef
Emery	5.8 fg	2.6 a	80.5 ab	4.7 ef	1.2 a	1.6 ef
Sullivan	6.1 de	2.4 cd	78.1 cd	6.7 b-d	1.2 ab	1.7 c
Walton	5.7 g	2.5 ab	79.7 ab	4.9 ef	1.2 a	1.9 a
Wynne	6.4 bc	2.6 ab	77.1 d	7.5 b	1.2 a	1.6 f
N14002oIJ	6.5 b	2.6 ab	77.5 d	7.1 bc	1.2 ab	1.6 ef
N14004oIJ	6.3 b-d	2.6 a	77.4 d	7.2 b	1.2 a	1.6 f
N14023ol	6.0 ef	2.2 ef	81.1 a	4.4 f	1.1 c	1.8 bc
N14027oIJ	6.5 b	2.2 ef	77.4 d	7.7 b	1.1 c	1.7 de
N15017ol	6.1 de	2.3 de	79.5 bc	5.4 d-f	1.2 b	1.8 b
N15039ol	6.0 ef	2.5 bc	79.4 bc	5.8 c-e	1.2 ab	1.6 ef
N15041ol	6.4 b	2.2 ef	78.1 cd	7.0 bc	1.1 c	1.7 cd
N15044oIF	6.5 b	2.2 f	77.2 d	7.8 b	1.1 c	1.7 c
Mean	6.4	2.4	76.8	7.9	1.2	1.7
LSD²	0.2	0.1	1.6	1.4	0.1	0.1

¹ Refer to page 3 for an explanation of the computations of these characters.

² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 31. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Three-year averages across all locations, (2018 – 2020)¹, (cont.)

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
Bailey	2.5 a	1.3 d	95.1 a	2.1 f	16.6 a	1.6 a	5.0 ab
Bailey II	2.3 b-f	1.3 d	80.7 b	14.6 cd	13.4 cd	0.5 bc	4.7 d-f
Emery	2.3 b-e	1.3 d	78.6 f	18.1 ab	13.2 cd	0.4 ef	4.8 c-e
Sullivan	2.3 bc	1.4 ab	80.1 b-d	14.2 cd	13.5 b-d	0.5 b-d	4.9 bc
Walton	2.6 a	1.5 a	78.5 f	18.1 ab	13.5 bc	0.4 ef	5.2 a
Wynne	2.3 bc	1.3 d	80.6 bc	11.4 e	13.8 b	0.5 b	4.8 cd
N14002oIJ	2.3 b	1.3 d	80.1 b-d	14.8 cd	13.8 b	0.5 bc	4.8 cd
N14004oIJ	2.3 b-d	1.3 cd	80.3 bc	13.0 de	13.8 b	0.5 b	4.8 cd
N14023oI	2.1 g	1.3 b-d	78.7 ef	19.3 a	12.8 e	0.3 f	4.6 f
N14027oIJ	2.1 fg	1.3 cd	81.2 b	14.4 cd	13.3 cd	0.6 b	4.6 f
N15017oI	2.3 bc	1.4 a-c	79.1 d-f	16.7 bc	13.3 cd	0.4 d-f	4.9 b-d
N15039oI	2.2 c-g	1.3 d	79.6 c-e	17.6 ab	13.2 de	0.4 c-e	4.7 d-f
N15041oI	2.2 e-g	1.3 b-d	80.6 bc	13.9 de	13.2 cd	0.5 bc	4.6 ef
N15044oIF	2.2 d-g	1.3 b-d	81.2 b	13.0 de	13.3 cd	0.6 b	4.6 ef
Mean	2.3	1.3	81.0	14.4	13.6	0.6	4.8
LSD²	0.1	0.1	1.1	2.6	0.3	0.1	0.2

¹ Refer to page 3 for an explanation of the computations of these characters.

² Minimum significant difference at P=0.05, based on the Fisher LSD test.

³ Lower iodine value indicates longer shelf life.

⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 32. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Rain Shelter Trial, Suffolk, VA 2020¹.

Variety	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
N14001	5.9 c	2.0 ab	79.8 a	6.4 c	1.0 ab	1.8 ef
N14002oIJ	6.4 bc	1.8 b-d	76.0 ab	9.3 bc	0.9 ab	2.1 a-c
N15066	6.5 bc	1.7 cd	75.5 ab	9.7 bc	0.9 b	2.2 a-c
N14023ol	5.8 c	1.6 d	79.2 a	6.7 c	0.9 b	2.4 a
N15053	6.8 bc	1.9 a-c	70.9 a-c	13.6 bc	1.0 ab	2.1 a-c
N15060	6.4 bc	1.7 cd	75.1 ab	9.8 bc	0.9 b	2.3 a
N16012	6.0 c	1.8 b-d	77.8 a	8.0 c	0.9 ab	2.1 a-f
N16021	6.5 bc	1.8 b-d	73.7 ab	11.0 bc	0.9 ab	2.2 a-d
N17036	5.9 c	2.2 a	80.1 a	5.5 c	1.1 a	1.9 c-f
N17037	5.8 c	1.9 a-c	79.1 a	6.3 c	0.9 ab	1.7 f
N17045	6.2 bc	1.7 cd	75.4 ab	10.1 bc	0.9 ab	2.3 ab
N17047	6.6 bc	1.6 cd	74.3 ab	10.7 bc	0.9 b	2.2 a-d
N04074FCT (ck)	7.7 ab	1.7 b-d	64.7 bc	19.3 ab	0.9 ab	1.9 d-f
SPT06-07 (ck)	8.6 a	1.8 b-d	59.0 c	23.9 a	1.0 ab	1.8 ef
Bailey II	5.9 c	1.9 b-d	80.1 a	6.0 c	1.0 ab	1.9 b-f
Walton	6.8 bc	1.9 a-c	71.0 a-c	13.1 bc	1.0 ab	2.1 a-c
Mean	6.5	1.8	74.5	10.6	0.9	2.1
LSD²	1.6	0.3	12.2	10.3	0.2	0.3

¹ Refer to page 3 for an explanation of the computations of these characters.² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 32. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Rain Shelter Trial, Suffolk, VA 2020¹, (cont.)

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
N14001	1.9 c	1.1 c	81.2 c	12.7 a-d	11.9 c	0.5 cd	4 c
N14002oIJ	2.2 a-c	1.3 a-c	83.1 c	8.9 a-g	12.6 bc	0.7 cd	4 a-c
N15066	2.3 a-c	1.3 a-c	83.4 bc	7.8 b-g	12.7 bc	0.8 cd	4 a-c
N14023ol	2.2 a-c	1.3 a-c	81.5 c	11.9 a-e	11.8 c	0.6 cd	4 a-c
N15053	2.4 a-c	1.3 a-c	86.2 bc	5.7 fg	13.4 a-c	1.0 a-c	5 a-c
N15060	2.4 a-c	1.4 a-c	83.3 bc	7.7 b-g	12.8 bc	0.8 cd	5 a-c
N16012	2.2 a-c	1.2 bc	82.4 c	9.8 a-f	12.1 bc	0.7 cd	4 bc
N16021	2.5 a-c	1.5 a	84.0 bc	6.7 d-g	13.2 a-c	0.8 b-d	5 ab
N17036	2.1 a-c	1.2 a-c	80.0 c	14.6 a	12.5 bc	0.4 d	4 a-c
N17037	2.8 a	1.4 a-c	80.2 c	13.5 a-c	12.9 bc	0.5 cd	5 a
N17045	2.1 bc	1.3 a-c	84.1 bc	7.5 c-g	12.3 bc	0.8 b-d	4 bc
N17047	2.4 a-c	1.4 a-c	84.1 bc	8.6 a-g	12.9 bc	0.8 b-d	5 a-c
N04074FCT (ck)	2.5 a-c	1.3 a-c	90.5 ab	4.1 fg	14.2 ab	1.3 ab	5 a-c
SPT06-07 (ck)	2.6 ab	1.3 a-c	93.5 a	3.0 g	15.3 a	1.5 a	5 ab
Bailey II	2.1 bc	1.2 c	80.8 c	13.6 ab	12.0 c	0.5 cd	4 bc
Walton	2.5 a-c	1.5 ab	85.4 bc	6.4 e-g	13.8 a-c	0.9 b-d	5 ab
Mean	2.3	1.3	84.0	8.9	12.9	0.8	4.6
LSD²	0.7	0.3	7.2	6.1	2.2	0.6	0.8

¹ Refer to page 3 for an explanation of the computations of these characters.

² Minimum significant difference at P=0.05, based on the Fisher LSD test.

³ Lower iodine value indicates longer shelf life.

⁴ Higher O/L ratio indicates longer shelf life.

Fatty Acid Results

Table 33. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Rain Shelter Trial, Suffolk, VA (2019-2020)¹ Two-year averages

	Palmitic C16:0	Stearic C18:0	Oleic C18:1	Linoleic C18:2	Arachidic C20:0	Eicosenoic C20:1
N14023	6.0 b	1.7 b	79.1 a	6.1 b	0.9 b	2.2 a
N17045	6.2 b	1.9 ab	77.5 a	7.6 b	1.0 ab	2.1 a
N17047	6.4 b	1.8 b	76.4 a	8.4 b	0.9 b	2.1 a
N04074FCT_(ck)	8.2 a	2.0 ab	60.9 b	22.0 a	1.0 ab	1.8 b
SPT06-07_(ck)	9.2 a	2.2 a	55.6 b	26.0 a	1.1 a	1.6 b
Mean	7.2	1.9	68.9	14.0	1.0	1.9
LSD	1.2	0.4	9.1	7.6	0.2	0.3

¹ Refer to page 3 for an explanation of the computations of these characters.

² Minimum significant difference at P=0.05, based on the Fisher's LSD test.

Fatty Acid Results

Table 33. Fatty Acid Composition, Iodine Values, Oleic/Linoleic (O/L) Ratio, % Total Saturated, Polyunsaturated/Saturated (P/S) Ratio, and % Total Long Chain Saturated. Rain Shelter Trial, Suffolk, VA (2019-2020)¹ Two-year averages (cont.)

Variety	Behenic C22:0	Lignoceric C24:0	Iodine Value ³	O/L ratio ⁴	% Total Saturated	P/S ratio	% Total Long Chain Saturated
N14023	2.4 b	1.4 a	80.4 b	13.0 a	12.6 b	0.5 b	4.8 a
N17045	2.4 b	1.4 a	81.5 b	11.6 a	12.7 b	0.6 b	4.7 a
N17047	2.5 ab	1.5 a	81.8 b	10.8 a	13.1 b	0.6 b	4.9 a
N04074FCT_(ck)	2.7 ab	1.4 a	91.9 a	3.2 b	15.3 a	1.4 a	5.2 a
SPT06-07_(ck)	2.9 a	1.4 a	94.1 a	2.4 b	16.8 a	1.5 a	5.4 a
Mean	2.6	1.4	86.0	8.2	14.1	0.9	5.0
LSD	0.5	0.2	5.6	4.8	2.1	0.4	0.8

¹ Refer to page 3 for an explanation of the computations of these characters.

² Minimum significant difference at P=0.05, based on the Fisher LSD test.

³ Lower iodine value indicates longer shelf life.

⁴ Higher O/L ratio indicates longer shelf life.